

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
28 June 2001 (28.06.2001)


PCT

(10) International Publication Number  
**WO 01/46408 A2**

- (51) International Patent Classification<sup>7</sup>: C12N 15/11, 15/86, 15/63, 5/10, 5/22, 1/21, A61K 39/21
- (21) International Application Number: PCT/US00/34985
- (22) International Filing Date:  
22 December 2000 (22.12.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
60/173,036 23 December 1999 (23.12.1999) US
- (71) Applicant (for all designated States except US): THE GOVERNMENT OF THE UNITED STATES OF AMERICA, as represented by THE SECRETARY, DEPARTMENT OF HEALTH AND HUMAN SERVICES [US/US]; Office of Technology Transfer, National Institutes of Health, 6011 Executive Blvd., Suite 325, Rockville, MD 20852 (US).
- (72) Inventor; and  
(75) Inventor/Applicant (for US only): PAVLAKIS, George, N. [US/US]; 9 Purdue Court, Rockville, MD 20850 (US).
- (74) Agents: FEILER, William, S. et al.; Morgan & Finnegan, L.L.P., 345 Park Avenue, New York, NY 10154 (US).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:  
— Without international search report and to be republished upon receipt of that report.

[Continued on next page]

(54) Title: MOLECULAR CLONES WITH MUTATED HIV GAG/POL, SIV GAG AND SIV ENV GENES



(57) Abstract: Nucleic acid constructs containing HIV-1 GAG/POL and SIV gag or SIV env genes which have been mutated to remove or reduce inhibitory/instability sequences are disclosed. Viral particles and host cells containing these constructs and/or viral particles are also disclosed. The exemplified constructs and viral particles of the invention may be useful in gene therapy for numerous disorders, including HIV infection, or as a vaccine for HIV-1 immunotherapy and immunoprophylaxis.

WO 01/46408 A2



---

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

**MOLECULAR CLONES WITH MUTATED  
HIV GAG/POL, SIV GAG AND SIV ENV GENES**

**I. TECHNICAL FIELD**

The invention relates to nucleic acids comprising mutated HIV-1 *gag/pol* and SIV *gag* gene sequences which are capable of being expressed independently of any SIV or HIV regulatory factors. The invention also relates to nucleic acids comprising a mutated SIV *env* gene sequence, which is capable of being expressed independently of any SIV or HIV regulatory factors. The preferred nucleic acids of the invention are capable of producing infectious viral particles.

The invention also relates to vectors, vector systems and host cells comprising the mutated HIV-1 *gag*, HIV-1 *pol* and/or SIV *gag* gene sequences. The invention also relates host cells comprising these nucleic acids and/or vectors or vector systems. The invention also relates to the use of these nucleic acids, vectors, vector systems and/or host cells for use in gene therapy or as vaccines.

**II. BACKGROUND**

Until recently, gene therapy protocols have often relied on vectors derived from retroviruses, such as murine leukemia virus (MLV). These vectors are useful because the genes they transduce are integrated into the genome of the target cells, a desirable feature for long-term expression. However, these retroviral vectors can only transduce dividing cells, which limits their use for *in vivo* gene transfer in nonproliferating cells, such as hepatocytes, myofibers, hematopoietic stem cells, and neurons.

Lentiviruses are a type of retrovirus that can infect both dividing and nondividing cells. They have proven extremely efficient at providing long-term gene expression (for up to 6 months) in a variety of nondividing cells (such as, neurons and macrophages) in animal models. See, e.g., Amado et al., *Science* 285:674-676 (July 1999). It has been proposed that the optimal gene transfer system would include a vector based on HIV, or other lentivirus, that can integrate

- 2 -

into the genome of nonproliferating cells. Because retroviruses integrate in the genome of the target cells, repeated transduction is unnecessary. Therefore, in contrast to an adenoviral vector capable of *in vivo* gene delivery, problems linked to the humoral response to injected viral antigens can be avoided. See, e.g., Naldini et al., *Science*, 272:263-267 (1996), p. 263.

HIV and other lentiviruses have a complex genome that, in addition to the essential structural genes (*env*, *gag*, and *pol*), contains regulatory (*tat* and *rev*) and accessory genes (*vpr*, *vif*, *vpu*, and *nef*). HIV has evolved to efficiently infect and express its genes in human cells, and is able to infect nondividing cells such as macrophages because its preintegration complex can traverse the intact membrane of the nucleus in the target cell. This complex contains, in addition to the viral DNA, the enzyme integrase, the product of the *vpr* gene, and a protein encoded by the *gag* gene called matrix. The matrix protein enables the preintegration complex to pass into the nucleus to access the host DNA. Lentiviruses cannot efficiently transduce truly quiescent cells (cells in the G<sub>0</sub> state). However, unlike murine retroviral vectors, in addition to being able to infect dividing cells, HIV-based vectors can achieve effective and sustained transduction and expression of therapeutic genes in nondividing cells, such as hematopoietic stem cells and in terminally differentiated cells such as neurons, retinal photoreceptors, muscle, and liver cells. See, e.g., Amado et al. (July 1999) and Klimatcheva et al., *Frontiers in Bioscience* 4:d481-496 (June 1999), and the references cited therein.

Although lentiviral vectors can be efficient gene delivery vehicles, there are safety concerns due to their origin. Therefore, the field has turned its attention to the development of vectors and production systems with built-in safety features to prevent the emergence of replication competent lentivirus (RCL). For example, in most laboratory applications, lentiviral vectors are generally created in a transient system in which a cell line is transfected with three separate constructs: a packaging construct, a transfer construct, and an envelope encoding construct. The packaging construct contains the elements necessary for vector packaging (except for *env*) and the enzymes required to generate vector particles. The transfer construct contains genetic *cis*-acting sequences necessary for the vector to infect the target cell and for transfer of the therapeutic (or reporter) gene. The lentivirus *env*

- 3 -

gene is generally deleted from the packaging construct and instead the envelope gene of a different virus is supplied in a third vector "the env-coding vector", although the lentiviruses *env* gene may be used if it is desired that the vector be intended to infect CD4<sup>+</sup>T cells. A commonly used envelope gene is that encoding the G glycoprotein of the vesicular stomatitis virus (VSV-G), which can infect a wide variety of cells and in addition confers stability to the particle and permits the vector to be concentrated to high titers (see, e.g., Naldini et al., Science 272:263-267 (1996) and Akkina et al. J. Virol. 70:2581 (1996). The use of three separate constructs and the absence of overlapping sequences between them minimizes the possibility of recombination during lentivirus (transfer) vector production. In addition, because no viral proteins are expressed by the lentiviral (transfer) vector itself, they do not trigger an effective immune response against cells expressing vector in animal models (a particular problem with vectors based on adenovirus). See, e.g., Amado et al., Science 285:674-676 (July 1999) and the references cited therein. See also Naldini et al. Science 272:263-267 (1996).

The initial packaging plasmids contained most HIV genes except for *env*. In an effort to improve safety, subsequent HIV vectors have been produced in which the packaging plasmid is devoid of all accessory genes. This process does not interfere with efficient vector production and significantly increases the safety of the system because potential RCLs lack the accessory genes necessary for efficient replication of HIV in humans. Although these vectors can transduce growth-arrested cell lines and neurons *in vivo*, they have been reported to not efficiently transduce macrophages. The accessory gene *vpr* is believed to be necessary for HIV infection of these cells using these HIV vectors. See, Zufferey et al., Nature Biotechnol. 15:871-875 (1997). In contrast, as discussed later herein, the HIV-based lentiviral vectors of the present invention do not need any HIV accessory genes in order to be able to infect human macrophages and the other cells tested.

The requirement of *vpr* or *vif* for efficient transduction of liver cells has also been reported. See, e.g., Kafri et al., Nature Genet. 17:314 (1997). These results indicate that the requirement of accessory genes for efficient lentivirus-mediated gene transfer is dependent on the type of cell chosen as target, suggesting

- 4 -

that future applications of lentiviral vectors may involve vector constructs with different accessory genes, as needed.

Zufferey et al., (1997) describe an HIV vector system in which the virulence genes, *env*, *vif*, *vpr*, *vpu*, and *nef* have been deleted. This multiply attenuated vector conserved the ability to transduce growth-arrested cells and monocyte-derived macrophages in culture, and could efficiently deliver genes in vivo into adult neurons. The packaging plasmids described Zufferey et al. (1997) and Naldini et al. (1996) encode Rev and Tat, in addition to Gag and Pol.

Lentiviral vectors engineered to become packaged into virions in the absence of the regulatory gene *tat* have also been described. See, e.g., Kim et al., J. Virol. 72:811-816 (1998) and Miyoshi et al. J. Virol. 72:8150-8157 (1998). In these vectors the *tat* gene has been removed from the packaging plasmid. Kim et al. state that *tat* is not necessary as long as the serial 5' LTR promoter is replaced with a strong constitutive promoter. It also has other advantages for HIV therapy. Replacement of the HIV-1 LTR with a constitutive HCMV promoter permits the use of anti-Tat molecules such as Tat transdominant mutants or Tat activation response element decoys as therapeutic agents, since they will not affect vector production. (see p. 814, col. 2). The removal of the *tat* gene eliminates an essential virulence factor that could contribute to a possible RCL. Kim et al. (1998) describe a vector system which does not contain *tat*, *vif*, *vpr*, *vpu* and *nef*. The preferred vector system includes the *rev* gene which, the authors state "with RRE, is required for efficient RNA handling in this system." (p. 811, col. 2). However, Kim et al. also constructed Rev independent constructs using CTE. Kim et al. state that the *rev*/RRE components could be removed by using a sequence such as the Mason-Pfizer monkey virus (MPMV) constitutive transport element (CTE), thereby eliminating all accessory proteins, but this leads to a significant reduction in titer.

Srinivasakumar et al., J. Virol. 71:5841-5848 (1997) describes the generation of stable HIV-1 packaging lines that constitutively express high levels of HIV-1 structural proteins in either a Rev-dependent or a Rev-independent fashion. These cell lines were used to assess gene transfer by using a HIV-1 vector expressing the hygromycin B resistance gene and to study the effects of Rev, Tat, and Nef on the vector titer. The Rev-independent cell lines were created by using

- 5 -

*gag-pol* and *env* expression vectors that contain the MPMV CTE. This article describes the construction of four plasmids, among others: CMV *gagpol*-RRE and pCMVenv, which require Rev coexpression for HIV-1 structural gene expression, and pCMV *gagpol*-CTE and pCMVenv-CTE, which do not. To create Rev-containing and Rev-independent packaging, cell lines, CMT3 cells were transfected with vectors expressing Gag, Gag-Pol, and Env, using a calcium phosphate transfection procedure.

By creating an HIV vector which contained the MPMV CTE (pTR167-CTE) and a packaging cell line which expressed the HIV structural proteins in a Rev-independent fashion, the authors were able to obtain a HIV vector system that functions completely without Rev. The titer of the vector obtained from this system was essentially the same as that obtained from a parallel system which contained Rev. The authors state that, in this context, the CTE seemed to substitute completely for Rev-RRE functions, similar to what was previously observed in transient-expression assays with Rev-dependent constructs. This is in contrast to situations where several rounds of HIV replication were measured. In those cases, titers from CTE-containing viruses were always reduced by at least 1 log unit compared to viruses utilizing Rev and the RRE. (See, Srinivasakumar et al., p. 5847).

The authors state that the advantages of having a HIV vector system that works in the absence of Rev opens the possibility of using it as a delivery vehicle for intracellular immunization against Rev function. Genes encoding Rev antagonists that have dramatic inhibitory effects on HIV replication, such as Rev M10 or RRE decoys, could be introduced into an HIV vector and put into cells normally infectable by HIV. Expression of the "anti-Rev" gene would be expected to dampen HIV infection. Any residual HIV replication should lead to activation of the vector LTR (by Tat) and create a vector-derived RNA that would be packaged by proteins derived from the infectious virus. In this scenario, the wild-type virus would act as a helper that may allow the spread of vector particles to previously nonimmunized cells. Because of the additional vector spread, it is likely that this type of scheme will be more effective in modulating HIV infection *in vivo* than one

- 6 -

based on traditional retrovirus vectors. The authors state that they are currently testing this approach in model systems. (See, Srinivasakumar et al., p. 5847).

Another development in the quest for a safe system is the so-called self-inactivating (SIN) vector. See, e.g., Yu et al., *Proc Natl Acad Sci USA* 83:3194-8 (1986) and Miyoshi et al., *J. Virol.* 72:8150 (1998). In Yu et al., a retrovirus-derived vector SIN vector was designed for the transduction of whole genes into mammalian cells. The SIN vector of Yu et al. contains a deletion of 299 base pairs in the 3' long terminal repeat (LTR), which includes sequences encoding the enhancer and promoter functions. When viruses derived from such vectors were used to infect NIH 3T3 cells, the deletion was transferred to the 5' LTR, resulting in the transcriptional inactivation of the provirus in the infected cell. Introduction of a hybrid gene (human metallothionein-promoted c-fos) into cells via a SIN vector was not associated with rearrangements and led to the formation of an authentic mRNA transcript, which in some cases was induced by cadmium. The vector described in Miyoshi et al. also contains a deletion the 3' (downstream) LTR. A sequence within the upstream LTR serves as a promoter under which the viral genome is expressed. The deletion introduced in the downstream LTR is transferred to the upstream LTR during reverse transcription. This deletion inactivates the LTR promoter and eliminates the production of vector RNA. The gene (or genes) to be transferred (e.g., a reporter or therapeutic gene) is expressed from an exogenous viral or cellular promoter that is inserted into the lentivirus vector. An important safety feature of SIN vectors is that inactivation of the promoter activity of the LTR reduces the possibility of insertional mutagenesis (of the transfer vector) into the host genome. In addition, because the expression of the (transfer) vector RNA is eliminated, the potential for RCL production in the target cell is further minimized. SIN vectors should be particularly useful in gene transfer experiments designed to study the regulated expression of genes in mammalian cells. Absence of enhancer and promoter sequences in both LTRs of the integrated provirus should also minimize the possibility of activating cellular oncogenes and may provide a safer alternative to be used in human gene therapy. Other modifications to enhance safety and specificity include the use of specific internal promoters that regulate gene expression, either temporally or with tissue or cell specificity.



- 7 -

Other strategies to improve safety in human studies would be to use nonhuman lentiviruses such as simian immunodeficiency virus, bovine immunodeficiency virus, or equine infectious anemia virus. Of these, vectors derived from the feline immunodeficiency virus have been engineered to efficiently transduce nondividing human cells. See, e.g., Poeschla et al., *Nature Med.* 4:354-357 (1998) and WO 99/15641. In addition, White et al., *J. Virol.* 73:2832-2840 (April 1999) described lentiviral vectors using human and simian immunodeficient virus elements in attempt to improve safety by reducing the likelihood of recombination between packaging constructs and transfer constructs.

The development of efficient packaging lines has proven challenging because expression of the VSV-G envelope and a number of HIV proteins is toxic to cells. Recently, a producer line has been designed in which the expression of packaging genes and VSV-G, and therefore the production of vector, can be turned on at will. Kafri et al., *J. Virol.* 73:576-584 (1999). The cell line can be expanded for scale-up vector production when the expression of toxic genes is turned off. This cell line produces high titer vector without generating RCL. Hematopoietic stem cells transduced with an HIV vector were transplanted into rhesus macaques as described by Donahue et al. *Blood* 92 (suppl. 1), abstract 4648.5 (1998) with at least a 14-month follow-up. At that time the procedure proved to be safe; all animals in the study have remained healthy without evidence of circulating HIV or vector. See, Amado et al., *Science* 285:674-676 (July 1999).

Many gene therapy protocols have been designed to correct a number of inherited metabolic, infectious, or malignant diseases using the hematopoietic stem cell. This cell has the capacity to self-renew and to differentiate into all of the mature cells of the blood and immune systems. Many diseases that affect these systems could potentially be treated by the stable introduction of therapeutic genes into stem cells. Recently, lentiviral vectors were shown to bypass the need for *ex vivo* stem cell stimulation (which is necessary when using murine retroviral vectors), by mediating efficient gene transfer into very primitive human stem cells that contributed to stable, long-term reconstitution of SCID mouse bone marrow with many hematopoietic lineages. See, e.g., Miyoshi et al., *Science* 283:682 (1999). Similarly, in a rhesus macaque model of autologous transplantation with

- 8 -

lentivirus-transduced stem cells, multilineage gene expression was found, suggesting transduction of an early blood cell progenitor under conditions of minimal stem cell stimulation, ordinarily insufficient for transduction with murine retroviruses. See, Donahue et al., Blood 92 (suppl. 1), abstract 4648.5 (1999) and Amado et al., Science 285:674-676 (July 1999).

In HIV infection, another advantage of lentiviral vectors designed against HIV is their potential to be mobilized by HIV in the infected patient, because the virus supplies all of the necessary elements for packaging of the vector. If these mobilized vectors contained the HIV envelope, they could efficiently transfer their genes (for example, genes custom-designed to confer resistance against HIV) into CD4<sup>+</sup> T cells, protecting them from subsequent HIV infection. Lentiviral vectors can also be designed to efficiently express their genes only in CD4<sup>+</sup> T cells that are infected with HIV (so called *tat*-inducible vectors). In these vectors, all HIV genes, including *tat* and *rev*, are ablated; cis-acting sequences required for integration, expression, and packaging are retained, and expression is dependent on the activity of the HIV LTR (which requires transactivation by Tat). It has been shown that in this system, vector expression is induced efficiently upon HIV infection. Moreover, in the absence of genes that confer resistance against HIV, stable integration of this vector in permissive cell lines resulted in inhibition of HIV replication. Although the mechanism of HIV inhibition has not been completely elucidated, preliminary results suggest that this vector competes with HIV at the level of reverse transcription. See, An et al., J. Virol., in press, and Amado et al., Science 285:674-676 (1999).

A number of other potential medical applications, where the modification of the genetic material of quiescent cells could result in the prevention or reversal of a disease process, are beginning to be explored. For example, the finding that lentiviral vectors can mediate stable and long-term gene transfer by direct injection of vector into the rat and mouse retina has lent support to the notion of gene therapy for the treatment of retinitis pigmentosa. This degenerative disease of the retina is characterized by photoreceptor cell death, resulting in a slow progression to blindness. Mutations in the cGMP phosphodiesterase  $\beta$  subunit

- 9 -

(PDE $\beta$ ) gene of rod photoreceptors lead to an autosomal recessive form of retinitis pigmentosa in humans, and in the *rd* mouse model of the disease. Previous studies have shown that adenovirus and adeno-associated virus-mediated PDEP subretinal gene transfer results in a delay in photoreceptor cell death. Using the *rd* mouse model, a recent study demonstrated that photoreceptors could be rescued in up to 50% of eyes injected with a lentivirus vector containing the murine PDE $\beta$  gene. In contrast with the short-term expression previously obtained with adenovirus vectors, PDE $\beta$  expression in this study persisted for at least 24 weeks. This finding points to the potential success of gene therapy in a disease that currently lacks effective treatment. See, Takahashi et al., J. Virol., 73:7812-7816 (Sept. 1999) and Amado et al. Science, 285:674-676 (1999).

In nature, the expression of *gag*, *pol*, and *env* of HIV-1 depends on the presence of the viral Rev protein. This dependence is, at least in part, due to the presence of negatively acting sequences (inhibitory or instability elements [INS]) located within unspliced and partially spliced mRNAs. The positive interaction of Rev with the Rev-responsive element [RRE] in these mRNAs counteracts the negative effects of the inhibitory sequences.

None of the above references teach or suggest that the *gag* and/or *pol* genes described therein may be replaced with the *gag* and/or *pol* genes in which the inhibitory/instability have been mutated to render their expression Rev-independent. Furthermore, there is no disclosure of the specific HIV-1 *gag/pol* or SIV *gag* mutated genes described herein.

The *gag/pol* clone of the invention was made using the method for eliminating inhibitory/instability regions from a gene as first described in U.S. patent application Serial No. 07/858,747, filed March 27, 1992, (inventors, G. Pavlakis and B. Felber) entitled "Method of Eliminating Inhibitory/Instability Regions from mRNA" and later described in a Continuation-in-Part ("CIP") application, filed as PCT application PCT/US93/02908 on March 29, 1993 and U.S. Patent Nos. 5,972,596 and 5,965,726. The disclosure of the CIP application was published as International Publication No. WO 93/20212 on October 14, 1993. (The disclosures of these patents and patent applications are specifically

- 10 -

incorporated by reference herein in their entirety.) The method was also described in Schwartz et al., J. Virol. 66:7176-7182 (1992).

Schneider et al., J. Virol. 71:4892-4903 (1997), extend the work described in the patent applications and in Schwartz et al. by identifying and characterizing additional INS within *gag*, *protease* and *pol* genes and mutating them in a similar manner. Schneider et al. disclose nucleic acid constructs which contain completely mutated HIV-1 *gag* genes, but only partially mutated HIV-1 *pol* genes.

Schneider et al. demonstrate that expression vectors containing an intact or nearly intact p55<sup>gag</sup> region allow the production of immature viral particles in mammalian cells in the absence of any other HIV proteins. The introduction of additional mutations in the *protease* region allowed efficient production of Gag/protease, which resulted in processing of the Pr55<sup>gag</sup> precursor and production of mature Gag particles with a lentivirus-like conical-core structure.

Schneider et al. disclose that Rev-independent expression vectors allow the efficient expression of Gag proteins in many cell lines that are not able to support efficient Rev-RRE-dependent rescue of these RNAs. Schneider et al. also disclose that *gag/pol* expression vectors may be important for vaccination approaches against HIV-1, since the *gag/pol* region is more conserved than is the *env* region and may be important for an effective immune response against HIV and for protection against infection. They also state that efficient HIV gene expression in many cells is also of interest for possible gene transfer experiments using lentiviral vectors in nondividing or slowly dividing cells, since HIV and the other lentiviruses are able to infect quiescent cells.

Pavlakakis et al., Natl Conf Hum Retroviruses Relat Infect (2nd). (1995), 91, state that Rev-independent Gag expression vectors were able to produce viral particles in human and mouse cells in the absence of any other HIV proteins, and that additional mutations in the *pol* region allowed the expression of the protease and the processing of the p55 gag precursor. Direct DNA injection of TAT and Rev independent Gag expression vectors in mouse muscle resulted in Gag expression detected by ELISA and in anti-gag antibody response. Several Rev-and Tat-independent Gag expression cassettes were inserted into retroviral vectors and

- 11 -

cell lines expressing Gag or Gag fragments that are dominant negative inhibitors of HIV-1 were constructed.

Shiver et al. (1996) describe the results of DNA vaccination of mice and non-human primates with mutated plasmid DNA encoding either mutated genes encoding HIV-1 gag (p55 gag) or env (gp120 or gp160). Both gag and env vaccine recipients exhibited antigen-specific cytotoxic and helper T lymphocyte (CTL, Th) responses. The results are stated to demonstrate that DNA vaccines elicited long-lived T cell responses in both mice and nonhuman primates that were disseminated throughout the lymphatics.

### **III. SUMMARY OF THE INVENTION**

The invention relates to nucleic acids comprising the nucleic acid sequence of the mutated HIV-1 gag/pol gene shown in Figure 1 (SEQUENCE ID NO:1) and vectors and vector systems comprising these nucleic acids.

The invention also relates to nucleic acids comprising the nucleic acid sequence of the mutated SIV gag gene shown in Figure 3 and vectors and vector systems comprising these nucleic acids.

The invention also relates to nucleic acids comprising the mutated SIV env gene shown in Figure 17 and vectors and vector systems comprising these nucleic acids.

The invention also relates to products produced by the nucleic acids, e.g., mRNA, protein, and infectious viral particles.

The invention also relates to compositions comprising these nucleic acids and/or their expression products.

The invention also relates to host cells comprising these nucleic acids, vector systems or viral particles.

The invention also relates to uses of these nucleic acids, vector systems, host cells, expression products, and/or compositions to produce mRNA, proteins, and/or infectious viral particles, and/or to induce antibodies and/or cytotoxic or helper T lymphocytes.

The invention also relates to the use of these nucleic acid constructs, vectors, vector systems and or host cells for use in immunotherapy and

- 12 -

immunoprophylaxis, e.g., as a vaccine, or in genetic therapy after expression, preferably in humans. The nucleic acid constructs of the invention can include or be incorporated into lentiviral vectors or other expression vectors or they may also be directly injected into tissue cells resulting in efficient expression of the encoded protein or protein fragment. These constructs may also be used for *in-vivo* or *in-vitro* gene replacement, e.g., by homologous recombination with a target gene *in-situ*.

#### IV. BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1. DNA sequence of a mutated HIV-1 *gag/pol* molecular clone (SEQUENCE ID NO:1). The gagpol terminator is located at positions 4305-4397 of SEQUENCE ID NO:1.

Fig. 2. Comparison of the sequence of the wild -type and mutated *pol* region in pCMVgagpolBNkan. Position #1 in the figure is position 2641 in plasmid pCMVgagpolBNkan. The comparison starts at position 1872 from the gag initiator ATG.

Fig. 3. DNA sequence of a mutated SIV *gag* molecular clone (SIVgagDX).

Fig. 4. Comparison of the mutated SIV *gag* DNA sequence in SIVgagDX with the wild type SIV sequence from Simian (macaque) immunodeficiency virus isolate 239, clone lambda siv 239-1 (GenBank accession No. M33262).

Fig. 5. Schematic diagram of some components of sample versions of a lentiviral system. BGH poly (A): bovine growth hormone poly (A) signal; MSD: mutated splice donor site;  $\psi$ : encapsidation signal; SD, splice donor site; SA, splice acceptor site; "X" indicates that the ATG codon of the partial *gag* gene sequence is mutated so that translation of this gene does not occur.

Fig. 6. Schematic diagram of the packaging construct pCMVgagpolBNkan.

Fig. 7. Schematic diagram of transfer construct 1: pmBCwCNluci. The packaging signal, the CMV promoter and the coding region for the luciferase gene are flanked by the 5' and 3' HIV-1 LTRs, which provide promoter and

- 13 -

polyadenylation signals, as indicated by the arrows. Three consecutive arrows indicate the U5, R, and U3 regions of the LTR, respectively. The transcribed portions of the LTRs are shown in black. Some restriction endonuclease cleavage sites are also indicated.

Fig. 8. Schematic diagram of transfer construct 1: pmBCmCNluci.

Symbols are as above.

Fig. 9. DNA sequence of packaging construct pCMVgagpolBNkan.

Fig. 10. DNA sequence of transfer construct 1: pmBCwCNluci.

Fig. 11. DNA sequence of transfer construct 1: pmBCmCNluci.

Figure 12:

Fig. 12. Nucleotide sequence of the region BssHII (711) to ClaI (830) in wild-type HIV-1 molecular clones HXB2 and NL4-3, and in the transfer constructs. The translation initiator signal for Gag protein (ATG) is underlined. pmBCwCNluci and pmBCmCNluci (transfer constructs 1 and 2) contain the sequence mBCwCN. Transfer construct 3 contains the sequence m2BCwCN. In contrast to the sequence mBCwCN, m2BCwCN has different mutations at the 5' splice site region and has an intact Gag ATG.

Fig. 13. Bar graph showing levels of gag protein that is released from cells upon transient transfection with pCMVgagpolBNkan (labeled pCMVBKkan in the figure).

Fig. 14. Bar graph showing reverse transcriptase activity from the Rev-independent gag-pol HIV-1 vector pCMVgagpolBNkan (labeled pCMVBKkan in the figure).

Fig. 15. Bar graphs showing the amount of luciferase per nanogram of p24 Gag protein detected in cells transduced with pCMVgagpolBNkan Rev-independent gag-HIV-1 based retroviral vectors. The results show that with pCMVgagpolBNkan Rev-independent gag-HIV-1 based retroviral vectors display high transduction efficiency in (A) 293 cells, (B) human lymphoid cells, (C) human myeloid cells (U937), as well as (D) non-dividing cells such as primary human macrophages.

Fig. 16. Schematic diagram of the SIV envelope encoding vector CMVkan/R-R-SIVenvCTE.

Fig. 17. DNA sequence of the SIV envelope encoding vector CMVkan/R-R-SIVenvCTE containing a mutated SIV env gene.

## V. MODES FOR CARRYING OUT THE INVENTION

It is to be understood that both the foregoing general description and the following detailed description are exemplary and explanatory only, and are not restrictive of the invention, as claimed. The accompanying drawings, which are incorporated in and constitute a part of the specification, illustrate an embodiment of the invention and, together with the description, serve to explain the principles of the invention.

One aspect of the invention comprises vectors that encode the Gag and/or Pol of HIV-1 in a Rev-independent manner. An example of such a vector which is described herein is the plasmid pCMVgagpolBNkan, which encodes the complete Gag and Pol of HIV-1 in a Rev-independent manner, and also contains a gene conferring kanamycin resistance. This plasmid is Tat and Rev-independent and was generated by eliminating the inhibitory/instability sequences present in the *gag/pol* mRNA without altering the amino acid sequence of the proteins coded by the genes.

The *gag/pol* clone of the invention is a DNA construct of the *gag/pol* region of HIV which has had the inhibitory/instability regions removed. The construct is expected to be useful as a component a new type of lentivirus vector for use in gene therapy or as a vaccine.

The *gag*, *pol* or *gag/pol* sequences of the invention can be highly expressed in human and other mammalian cells in the absence of any other regulatory and structural protein of HIV, including Rev. When the *gag/pol* sequences are combined with a sequence encoding an envelope protein, such as the VSV G protein or the HIV envelope protein (e.g., in the same vector or in another expression vector), infectious virus is produced after transfection into human cells. When a gene encoding a non-HIV envelope protein is used, for example, in the presence of the HIV *gag/pol* gene, the virus particles produced would contain only the HIV proteins Gag and Pol.



- 15 -

Lentiviral vectors or vector systems based on the *gag*, *pol* or *gag/pol* sequences of this invention, as exemplified by the Rev-independent pCMVgagpol BNkan construct described herein, may be used for gene therapy *in vivo* (e.g., parenteral inoculation of high titer vector) or *ex vivo* (e.g., *in vitro* transduction of patient's cells followed by reinfusion into the patient of the transduced cells). These procedures are been already used in different approved gene therapy protocols.

The HIV *gag/pol* clone and SIV *gag* clone of the invention were made using the method for eliminating inhibitory/instability regions from a gene as described in U.S. patent application Serial No. 07/858,747, filed March 27, 1992, and also in related U.S. Patent Nos. 5,972,596 and 5,965,726, which are incorporated by reference herein. This method does not require the identification of the exact location or knowledge of the mechanism of function of the INS.

Generally, the mutations are such that the amino acid sequence encoded by the mRNA is unchanged, although conservative and non-conservative amino acid substitutions are also envisioned where the protein encoded by the mutated gene is substantially similar to the protein encoded by the non-mutated gene. The mutated genes can be synthetic (e.g., synthesized by chemical synthesis), semi-synthetic (e.g., a combination of genomic DNA, cDNA, or PCR amplified DNA and synthetic DNA), or recombinantly produced. The genes also may optionally not contain introns. The nucleic acids of the invention may also contain Rev-independent fragments of these genes which retain the desired function (e.g., for antigenicity of Gag or Pol, particle formation (Gag) or enzymatic activity (Pol)), or they may also contain Rev-independent variants which have been mutated so that the encoded protein loses a function that is unwanted in certain circumstances. In the latter case, for example, the gene may be modified to encode mutations (at the amino acid level) in the active site of reverse transcriptase or integrase proteins to prevent reverse transcription or integration. Rev-independent fragments of the *gag* gene are described in U.S. patent application Serial No. 07/858,747, filed March 27, 1992, and also in related U.S. Patent Nos. 5,972,596 and 5,965,726, which are incorporated by reference herein.

In addition to being capable of producing HIV Gag and Pol proteins in the absence of Rev regulatory protein in a cell *in vivo*, the HIV *gag/pol* clone and

- 16 -

SIV *gag* clone of the invention are also capable of producing HIV Gag and Pol proteins in the absence of any added cis acting transport element, such as CTE or CTE-like elements (collectively referred herein as RNA Transport Elements (RTE)). Experiments indicate that the mutated vectors of the invention for SIV *gag* are far superior to those adding CTE (see Qiu et al., J Virol. 73:9145-52 (1999)).

The expression of the proteins encoded by these vectors after transfection into human cells may be monitored at both the level of RNA and protein production. RNA levels are quantitated by methods known in the art, e.g., Northern blots, S1 mapping or PCR methods. Protein levels may also be quantitated by methods known in the art, e.g., western blot or ELISA or fluorescent detection methods. A fast non-radioactive ELISA protocol can be used to detect gag protein (DUPONT or COULTER gag antigen capture assay).

At least three types of lentiviral vectors based on the *gag/pol* genes of the invention for use in gene therapy and/or as a vaccine are envisioned, i.e., lentiviral vectors having

- a) no round of replication (i.e., a zero replication system)
- b) one round of replication
- c) a fully replicating system

For a system with no round of replication, a *gag/pol* gene, or separate *gag* and *pol* genes, or fragments of these genes, expressed using appropriate transcription units, e.g., a CMV promoter and a BGH poly (A) site. This will allow expression of the *gag/pol* unit (or *gag* or *pol* or fragment(s) thereof) for vaccine purposes. This expression can be accomplished without the production of any functional retroviral enzymes, provided that the appropriate mutation(s), e.g., a missense mutation, are introduced. In a zero replication system, a virus stock will be administered to the cells or animals of interest. For example, if one creates and uses a virus stock with the exemplified system using the packaging vector PCMVgagpolBNKan, the transfer construct pmBCwCNI<sub>luc</sub>i or pmBCmCNI<sub>luc</sub>i, and the envelope containing vector pHCMV-G, one obtains a zero replication system. The virus particles produced by such system can infect cells, and the reverse transcribed transfer construct DNA will go into the nucleus but, because the coding regions for viral structural proteins are not present, there will be no virus expression

- 17 -

and replication (0 rounds). If one transfects cells *in vivo* with the same 3 DNAs, they will go to the nucleus, express viral proteins, make infectious virus particles and go out and infect another cell or cells (1 round). Since *in vivo* delivery of three plasmids may result in lower expression, at least two different embodiments are envisioned. In the first, two plasmids may be used, e.g., MV1 shown in Fig. 5 and an envelope expression plasmid such as pHCMV-G. Other plasmids encoding functional envelopes from HIV, SIV, or other retroviruses can also be used. Transfection by the two plasmids results in infectious virus that can infect and integrate into new cells (1 round). The infected cells produce gagpol but virus propagation is not possible in the absence of env.

For a system with one round of replication, at least two additional embodiments are envisioned. In the first method, a combination of the genes, e.g., a *gag/pol* gene, an *env* encoding gene and, preferably, a gene encoding a reporter protein or other polynucleotide or protein of interest, are delivered into the cells of interest *in vivo*. As discussed above for the exemplified system, if one transfects cells *in vivo* with the same 3 DNAs, they will go to the nucleus, express viral proteins, make infectious virus particles, be released and infect another cell or cells (1 round).

In another embodiment, the same result (i.e., only one round of replication) can be obtained by using transfer vectors that have deletions in the 3' LTR and in which a heterologous-promoter (e.g., the CMV-promoter, or inducible promoter, or tissue-specific promoter), is used in place of the 3' LTR promoter. The mutations in the 3' LTR making it inactive upon reverse transcription and integration. This is because the integrated provirus derives both its 5' LTR and its 3' LTR from the 3' LTR of the starting (transfer) construct. (This is a well-known property of all retroviruses and has been used to make self-inactivating vectors (SIN)). There are several reasons one may want to inactivate the incoming LTR promoter, one of which is to use a different tissue specific or regulated promoter for expression of a gene of interest in the integrated provirus. Note that, with SIN vectors, if one uses a viral stock made *in vitro* after transfection into cells and collection of infectious virus, there will be no round of replication. If one transfects cells with the DNAs *in vivo*, there will be one round of replication. If functional

- 18 -

*gag*, *pol*, or *env* are not included in the DNA mix, there will not be any infection at all (i.e., infectious viruses will not be made).

A fully replicating Rev-independent system has not been constructed yet, although it is expected that a functional system can be constructed using Rev-independent *gag/pol* and *env* sequences. If desired, extra posttranscriptional control elements such as the CTE element, which can replace Rev and give infectious virus (see e.g., Zolotukhin et al., J.Virol.68:944-7952 (1994)) are included. The fully replicating system should be in one piece, containing the LTR, packaging signal, *gag/pol*, splice site, *env*, *tat*, one or more CTE or CTE-like elements (if desired for optimal results), and LTR. Tat is thought to be required in this construct, at least in non-permissive cells. Such a system is depicted in Figure 5, (construct MV2). In this system, a cell or animal of interest (preferably human) would be infected with virus stock that then propagates. CTE or CTE-like elements (depicted in construct MV2 as RTE (RNA Transport Elements)) are desirable since they have been shown to improve expression, and since many retroviruses require the presence of posttranscriptional control elements. There are several types of CTE and CTE-like elements, and these elements appear to work via a different pathway from the Rev-RRE pathway. See, e.g., Tabernero et al., J Virol. 71:95-101 (1997). See also, Pavlakis and Nappi, PCT/US99/11082, filed May 22, 1999, published as WO 99/61596 on December 2, 1999 (and incorporated herein by reference), which describes a new type of post-transcriptional control element that is able to replace CTE and HIV RRE/Rev. The Pavlakis-Nappi element does not work in the same way as CTE and does not have any sequence or structure homology.

In a preferred embodiment, a lentiviral system of the invention comprises the following three components:

1. a packaging vector containing nucleic acid sequences encoding the elements necessary for vector packaging such as structural proteins (except for HIV *env*) and the enzymes required to generate vector particles, the packaging vector comprising at least a mutated HIV or SIV *gag/pol* gene of the invention;

- 19 -

2. a transfer vector containing genetic cis-acting sequences necessary for the vector to infect the target cell and for transfer of the therapeutic or reporter or other gene(s) of interest, the transfer vector comprising the encapsidation signal and the gene(s) of interest or a cloning site for inserting the gene(s) of interest;

and

3. a vector containing sequences encoding an element necessary for targeting the viral particle to the intended recipient cell, preferably the gene encoding the G glycoprotein of the vesicular stomatis virus (VSV-G) or amphotrophic MuLV or lentiviral *envs*.

Using the CMV promoter or other strong, high efficiency, promoter instead of the HIV-1 LTR promoter in the packaging vector, high expression of *gag*, *pol*, or *gag/pol* can be achieved in the total absence of any other viral protein. The exchange of the HIV-1 LTR promoter with other promoters is beneficial in the packaging vector or other vectors if constitutive expression is desirable and also for expression in other mammalian cells, such as mouse cells, in which the HIV-1 promoter is weak. Vectors containing the sequences of the invention can be used for the Rev independent production of HIV-1 Gag/Pol, HIV-1 Gag, HIV-1 Pol, and SIV Gag proteins. In certain embodiments, the presence of heterologous promoters will also be desired in the transfer vector and the envelope encoding vector, when such vectors are used.

The gene(s) of interest are chosen according to the effect sought to be achieved. For gene therapy purposes there will be at least one therapeutic gene encoding a gene product which is active against the condition it is desired to treat or prevent. Alternatively or additionally, there may be a gene which acts as a marker by encoding a detectable product. Therapeutic genes may encode, for example, an anti-sense RNA, a ribozyme, a transdominant negative mutant of a target protein, a toxin, a conditional toxin, an antigen that induces antibodies or helper T-cells or cytotoxic T-cells, a single chain antibody or a tumor suppresser protein. See, e.g., WO 98/17816.

- 20 -

An even more extensive list of genes of interest for use in lentiviral vectors is described, e.g., in WO 99/04026 on page 10, line 20 to page 12, line 7. Table 2 of Klimatcheva et al. (1999) also provides a list of disorders and target cells for gene therapy, as well as a number of lentiviral vectors used by others. This list includes genetic/metabolic deficiencies, viral infection and cancer. Inherited genetic defects such as adenosine deaminase deficiency, familial hypercholesterolemia, cystic fibrosis, mucopolysaccharidosis type VII, types I and II diabetes, classical phenylketonuria and Gaucher disease are diseases which are listed as being possible to overcome by lentiviral vector-mediated gene therapy because they constitute single-gene deficiencies for which the involved genes are known. Viral diseases are also listed as constituting appropriate targets for lentiviral gene delivery. In particular, a number of gene therapy approaches have been proposed for the treatment of HIV infection and, for some of these strategies, phase I studies have recently begun in humans. The article states that preliminary studies have dealt with defective murine oncoviruses for delivery of anti-sense RNAs, ribozymes and trans-dominant proteins against HIV replication.

In any of the vectors, but preferably in the transfer vector, an inserted gene could have an internal ribosomal entry site (IRES), e.g., from picornaviral RNA. An IRES will be used in circumstances that one wants to express two proteins from the same promoter. For example one protein of interest and a marker gene, e.g., green fluorescent protein (GFP) or a marker gene and a drug resistance gene (e.g. the firefly luciferase gene and neomycin phosphotransferase gene) as described on p. 58 of WO 99/04026, for example. Using an IRES the expression of the two proteins is coordinated. A further gene or genes may also be present under the control of a separate promoter. Such a gene may encode for example a selectable marker, or a further therapeutic agent which may be among the therapeutic agents listed above. Expression of this gene may be constitutive; in the case of a selectable marker this may be useful for selecting successfully transfected packaging cells, or packaging cells which are producing particularly high titers of the retroviral vector particles. Alternatively or additionally, the selectable marker may be useful for selecting cells which have been successfully infected with the lentiviral vector and have the provirus integrated into their own genome.

- 21 -

One way of performing gene therapy is to extract cells from a patient, infect the extracted cells with a lentiviral vector and reintroduce the cells back into the patient. A selectable marker may be used to provide a means for enriching for infected or transduced cells or positively selecting for only those cells which have been infected or transduced, before reintroducing the cells into the patient. This procedure may increase the chances of success of the therapy. Selectable markers may be for instance drug resistance genes, metabolic enzyme genes, or any other selectable markers known in the art. Typical selection genes encode proteins that confer resistance to antibiotics and other toxic substances, e.g., histidinol, puromycin, hygromycin, neomycin, methotrexate etc. and cell surface markers.

However, it will be evident that for many gene therapy applications of lentiviral vectors, selection for expression of a marker gene may not be possible or necessary. Indeed expression of a selection marker, while convenient for *in vitro* studies, could be deleterious *in vivo* because of the inappropriate induction of cytotoxic T lymphocytes (CTLs) directed against the foreign marker protein. Also, it is possible that for *in vivo* applications, vectors without any internal promoters will be preferable. The presence of internal promoters can affect for example the transduction titres obtainable from a packaging cell line and the stability of the integrated vector. Thus, single transcription unit vectors, which may be bi-cistronic or poly-cistronic, coding for one or two or more therapeutic genes, may be the preferred vector designed for use *in vivo*. See, e.g., WO 98/17816.

Suitable host or producer cells for use in the invention are well known in the art. Many lentiviruses have already been split into replication defective genomes and packaging components. For those which have not the technology is available for doing so. The producer cell encodes the viral components not encoded by the vector genome such as the Gag, Pol and Env proteins. The *gag*, *pol* and *env* genes may be introduced into the producer cell transiently, or may be stably integrated into the cell genome to give a packaging cell line. The lentiviral vector genome is then introduced into the packaging cell line by transfection or transduction to create a stable cell line that has all of the DNA sequences required to produce a lentiviral vector particle. Another approach is to introduce the different

- 22 -

DNA sequences that are required to produce lentiviral vector particle, e.g., the *env* coding construct, the *gag-pol* coding construct and the transfer construct into the cell simultaneously by transient triple transfection.

Target cells identified by Klimatcheva et al. (1999), and the references cited therein, include airway epithelial cells for cystic fibrosis; retinal photoreceptor cells for retinitis pigmentosa; progenitors for red blood cells, macrophages, and lymphocytes for hematopoietic disorders, sickle cell anemia,  $\beta$ -thalassemia, lysosomal storage disorders, mucopolysaccharidoses, and severe combined immunodeficiency syndrome; bone marrow cells and macrophages for Gaucher's disease; liver cells for familial hypercholesterolaemia; T-lymphocytes and macrophages for HIV infection; brain tissue, neurons, and glial cells for neurodegenerative diseases such as Parkinson's and Alzheimer's diseases; endothelial cells and cardiac myocytes for cardiovascular diseases; and cancer cells in various tissues (e.g. liver or brain) for cancer. Target cells for other diseases would be apparent to one of skill in the art.

Vaccines and pharmaceutical compositions comprising at least one of the nucleic acid sequences, vectors, vector systems, or transduced or transfected host cells of the invention and a physiologically acceptable carrier are also part of the invention.

As used herein, the term "transduction" generally refers to the transfer of genetic material into the host via infection, e.g., in this case by the lentiviral vector. The term "transfection" generally refers to the transfer of isolated genetic material into cells via the use of specific transfection agents (e.g., calcium phosphate, DEAE Dextran, lipid formulations, gold particles, and other microparticles) that cross the cytoplasmic membrane and deliver some of the genetic material into the cell nucleus.

Systems similar to those described herein can be produced using elements of lentiviruses in addition to the HIV and/or SIV genes described herein.

#### Pharmaceutical Compositions

The pharmaceutical compositions of the invention contain a pharmaceutically and/or therapeutically effective amount of at least one nucleic acid



- 23 -

construct, vector, vector system, viral particle/virus stock, or host cell (i.e., agents) of the invention. In one embodiment of the invention, the effective amount of an agent of the invention per unit dose is an amount sufficient to cause the detectable expression of the gene of interest. In another embodiment of the invention, the effective amount of agent per unit dose is an amount sufficient to prevent, treat or protect against deleterious effects (including severity, duration, or extent of symptoms) of the condition being treated. The effective amount of agent per unit dose depends, among other things, on the species of mammal inoculated, the body weight of the mammal and the chosen inoculation regimen, as is well known in the art. The dosage of the therapeutic agents which will be most suitable for prophylaxis or treatment will also vary with the form of administration, the particular agent chosen and the physiological characteristics of the particular patient under treatment. The dose is administered at least once. Subsequent doses may be administered as indicated.

To monitor the response of individuals administered the compositions of the invention, mRNA or protein expression levels may be determined. In many instances it will be sufficient to assess the expression level in serum or plasma obtained from such an individual. Decisions as to whether to administer another dose or to change the amount of the composition administered to the individual may be at least partially based on the expression levels.

The term "unit dose" as it pertains to the inocula refers to physically discrete units suitable as unitary dosages for mammals, each unit containing a predetermined quantity of active material (e.g., nucleic acid, virus stock or host cell) calculated to produce the desired effect in association with the required diluent. The titers of the virus stocks to be administered to a cell or animal will depend on the application and on type of delivery (e.g., *in vivo* or *ex vivo*). The virus stocks can be concentrated using methods such as centrifugation. The titers to be administered *ex vivo* are preferably in the range of 0.001 to 1 infectious unit /cell. Another method of generating viral stocks is to cocultivate stable cell lines expressing the virus with the target cells. This method has been used to achieve better results when using traditional retroviral vectors because the cells can be infected over a longer period of time and they have the chance to be infected with multiple copies of the vector.

- 24 -

For *in vivo* administration of nucleic acid constructs, vectors, vector systems, virus stocks, or cells which have been transduced or transfected *ex vivo*, the dose is to be determined by dose escalation, with the upper dose being limited by the onset of unacceptable adverse effects. Preliminary starting doses may be extrapolated from experiments using lentiviral vectors in animal models, by methods known in the art, or may be extrapolated from comparisons with known retroviral (e.g., adenoviral) doses. Generally, small dosages will be used initially and, if necessary, will be increased by small increments until the optimum effect under the circumstances is reached. Exemplary dosages are within the range of  $10^8$  up to approximately  $5 \times 10^{15}$  particles.

Inocula are typically prepared as a solution in a physiologically acceptable carrier such as saline, phosphate-buffered saline and the like to form an aqueous pharmaceutical composition.

The agents of the invention are generally administered with a physiologically acceptable carrier or vehicle therefor. A physiologically acceptable carrier is one that does not cause an adverse physical reaction upon administration and one in which the nucleic acids are sufficiently soluble to retain their activity to deliver a pharmaceutically or therapeutically effective amount of the compound. The pharmaceutically or therapeutically effective amount and method of administration of an agent of the invention may vary based on the individual patient, the indication being treated and other criteria evident to one of ordinary skill in the art. A therapeutically effective amount of a nucleic acid of the invention is one sufficient to prevent, or attenuate the severity, extent or duration of the deleterious effects of the condition being treated without causing significant adverse side effects. The route(s) of administration useful in a particular application are apparent to one of ordinary skill in the art.

Routes of administration of the agents of the invention include, but are not limited to, parenteral, and direct injection into an affected site. Parenteral routes of administration include but are not limited to intravenous, intramuscular, intraperitoneal and subcutaneous. The route of administration of the agents of the invention is typically parenteral and is preferably into the bone marrow, into the CSF intramuscular, subcutaneous, intradermal, intraocular, intracranial, intranasal,

- 25 -

and the like. See, e.g., WO 99/04026 for examples of formulations and routes of administration.

The present invention includes compositions of the agents described above, suitable for parenteral administration including, but not limited to, pharmaceutically acceptable sterile isotonic solutions. Such solutions include, but are not limited to, saline and phosphate buffered saline for nasal, intravenous, intramuscular, intraperitoneal, subcutaneous or direct injection into a joint or other area.

In providing the agents of the present invention to a recipient mammal, preferably a human, the dosage administered will vary depending upon such factors as the mammal's age, weight, height, sex, general medical condition, previous medical history and the like.

The administration of the pharmaceutical compositions of the invention may be for either "prophylactic" or "therapeutic" purpose. When provided prophylactically, the compositions are provided in advance of any symptom. The prophylactic administration of the composition serves to prevent or ameliorate any subsequent deleterious effects (including severity, duration, or extent of symptoms) of the condition being treated. When provided therapeutically, the composition is provided at (or shortly after) the onset of a symptom of the condition being treated.

For all therapeutic, prophylactic and diagnostic uses, one or more of the agents of the invention, as well as antibodies and other necessary reagents and appropriate devices and accessories, may be provided in kit form so as to be readily available and easily used.

Where immunoassays are involved, such kits may contain a solid support, such as a membrane (e.g., nitrocellulose), a bead, sphere, test tube, rod, and so forth, to which a receptor such as an antibody specific for the target molecule will bind. Such kits can also include a second receptor, such as a labeled antibody. Such kits can be used for sandwich assays to detect toxins. Kits for competitive assays are also envisioned.

## VI. INDUSTRIAL APPLICABILITY

Mutated genes of this invention can be expressed in the native host cell or organism or in a different cell or organism. The mutated genes can be introduced into a vector such as a plasmid, cosmid, phage, virus or mini-chromosome and inserted into a host cell or organism by methods well known in the art. In general, the mutated genes or constructs containing these mutated genes can be utilized in any cell, either eukaryotic or prokaryotic, including mammalian cells (e.g., human (e.g., HeLa), monkey (e.g., Cos), rabbit (e.g., rabbit reticulocytes), rat, hamster (e.g., CHO and baby hamster kidney cells) or mouse cells (e.g., L cells), plant cells, yeast cells, insect cells or bacterial cells (e.g., *E. coli*). The vectors which can be utilized to clone and/or express these mutated genes are the vectors which are capable of replicating and/or expressing the mutated genes in the host cell in which the mutated genes are desired to be replicated and/or expressed. See, e.g., F. Ausubel et al., Current Protocols in Molecular Biology, Greene Publishing Associates and Wiley-Interscience (1992) and Sambrook et al. (1989) for examples of appropriate vectors for various types of host cells. The native promoters for such genes can be replaced with strong promoters compatible with the host into which the gene is inserted. These promoters may be inducible. The host cells containing these mutated genes can be used to express large amounts of the protein useful in enzyme preparations, pharmaceuticals, diagnostic reagents, vaccines and therapeutics.

Mutated genes or constructs containing the mutated genes may also be used for in-vivo or in-vitro gene therapy. For example, a mutated gene of the invention will produce an mRNA in situ to ultimately increase the amount of protein expressed. Such gene include viral genes and/or cellular genes. Such a mutated gene is expected to be useful, for example, in the development of a vaccine and/or genetic therapy.

The constructs and/or proteins made by using constructs encoding the mutated gag, env, and pol genes could be used, for example, in the production of diagnostic reagents, vaccines and therapies for AIDS and AIDS related diseases. The inhibitory/instability elements in the HIV-1 gag gene may be involved in the

- 27 -

establishment of a state of low virus production in the host. HIV-1 and the other lentiviruses cause chronic active infections that are not cleared by the immune system. It is possible that complete removal of the inhibitory/instability sequence elements from the lentiviral genome would result in constitutive expression. This could prevent the virus from establishing a latent infection and escaping immune system surveillance. The success in increasing expression of the entire *gag/pol* gene by eliminating the inhibitory sequence element suggests that one could produce lentiviruses without any negative elements. Such lentiviruses could provide a novel approach towards attenuated vaccines.

For example, vectors expressing high levels of Gag can be used in immunotherapy and immunoprophylaxis, after expression in humans. Such vectors include retroviral vectors and also include direct injection of DNA into muscle cells or other receptive cells, resulting in the efficient expression of gag, using the technology described, for example, in Wolff et al., Science 247:1465-1468 (1990), Wolff et al., Human Molecular Genetics 1(6):363-369 (1992) and Ulmer et al., Science 259:1745-1749 (1993). Further, the gag constructs could be used in transdominant inhibition of HIV expression after the introduction into humans. For this application, for example, appropriate vectors or DNA molecules expressing high levels of p55<sup>gag</sup> or p37<sup>gag</sup> would be modified to generate transdominant gag mutants, as described, for example, in Trono et al., Cell 59:113-120 (1989). The vectors would be introduced into humans, resulting in the inhibition of HIV production due to the combined mechanisms of gag transdominant inhibition and of immunostimulation by the produced gag protein. In addition, the gag constructs of the invention could be used in the generation of new retroviral vectors based on the expression of lentiviral gag proteins. Lentiviruses have unique characteristics that may allow the targeting and efficient infection of non-dividing cells. Similar applications are expected for vectors expressing high levels of env.

Identification of similar inhibitory/instability elements in SIV indicates that this virus is a convenient model to test these hypotheses. SIV similarly modified could be used in place of HIV in an effort to further minimize the possibility of rearrangement events that would lead to the generation of infectious HIV.

- 28 -

The following examples illustrate certain embodiments of the present invention, but should not be construed as limiting its scope in any way. Certain modifications and variations will be apparent to those skilled in the art from the teachings of the foregoing disclosure and the following examples, and these are intended to be encompassed by the spirit and scope of the invention.

### EXAMPLE 1

#### Rev-Independent HIV-1 Gag/Pol Molecular Clone

Figure 1 shows the DNA sequence of a Rev-independent HIV-1 *gag/pol* molecular clone. This DNA sequence shown encodes the complete Gag and Pol of HIV-1 and can be expressed in a Rev-independent manner when operably linked to a promoter. The Rev-independent *gag* sequence was described in U.S. Patent Nos. 5,972,596 and 5,965,726 and the Rev-independent *pol* sequence was generated by eliminating the inhibitory/instability sequences using the methods described in U.S. Patent Nos. 5,972,596 and 5,965,726. Others have reportedly made Rev independent *gag* sequences by optimizing codon usage for human cells (see, e.g., WO 98/34640).

Figure 2 shows an alignment of the sequence of the wild - type and mutated *pol* region in pCMVgagpolBNkan. Position #1 in the figure is position 2641 in plasmid pCMVgagpolBNkan.

The elimination of INS in *gag*, *pol* and *env* regions allows the expression of high levels of authentic HIV-1 structural proteins in the absence of the Rev regulatory factor of HIV-1.

### EXAMPLE 2

#### Rev-Independent SIV Gag Molecular Clone

Figure 3 shows the DNA sequence of a Rev-independent SIV *gag* molecular clone, SIVgagDX. Figure 4 shows the comparison of wild type (WT) and mutant (SIVgagDX) sequences. The wild type SIV sequence is from Simian (macaque) immunodeficiency virus isolate 239, clone lambda siv 239-1 (GenBank accession No. M33262).

- 29 -

### EXAMPLE 3

#### Rev-Independent SIV Env Molecular Clone

Figure 16 shows a schematic diagram, and figure 17 shows the DNA sequence, of the "env-coding" vector CMVkan/R-R-SIVenvCTE, which is an example of a vector comprising a mutated lentiviral *env* gene sequence which is capable of being expressed independently of any SIV or HIV regulatory factors. "CMV" denotes the cytomegalovirus promoter; "SRV-CTE" denotes the constitutive transport element (CTE) of Simian Retrovirus Type 1; "all-STOP" denotes a sequence providing translational stops in all three reading frames; "BGH terminator" denotes the bovine growth hormone polyadenylation signal. Other posttranscriptional control elements can be used instead of the indicated SRV-CTE, for example the one described by Pavlakis and Nappi, PCT/US99/11082, filed May 22, 1999, which was published as WO 99/61596 on December 2, 1999 (and which is incorporated herein by reference).

As mentioned previously above, such a vector encoding a lentiviral *env* gene may be used if it is desired that the vector infect CD4<sup>+</sup>T cells. Also as mentioned previously above, the CTE element (i.e., the SRV-CTE element in the case of vector CMVkan/R-R-SIVenvCTE), can be replaced with another post-transcriptional control element, such as the Pavlakis-Nappi element, that is able to replace CTE and HIV RRE/Rev. See Pavlakis and Nappi, PCT/US99/11082, filed May 22, 1999, which was published as WO 99/61596 on December 2, 1999 (and which is incorporated herein by reference).

### EXAMPLE 4

#### Lentiviral Vector System

Figure 5 is a schematic of some of the components of a preliminary version of the Rev-independent lentiviral vector system exemplified herein, including a packaging construct and three different transfer vectors which may be used. In the lentiviral system exemplified herein, the packaging construct also contains the gene for kanamycin resistance. The lentiviral system exemplified herein also contains the vector pHCMV-G, which is shown in Figure 5.

- 30 -

In the packaging construct shown in Figure 5, "CMV" denotes the cytomegalovirus promoter, "Gag" denotes the gag gene, which generates components of the virion core, "Pro" denotes "protease" "RT" denotes "reverse transcriptase," "Int" denotes "integrase" and "BGH poly (A)" denotes the bovine growth hormone polyadenylation signal. The protease, reverse transcriptase, and integrase genes comprise the "pol" gene. In transfer construct 1, "LTR" denotes the HIV "long terminal repeat", which contains a HIV promoter; "mSD" denotes "mutated splice donor site," which is present in the construct so that splicing of the RNA transcript does not occur; "ψ" denotes the encapsidation signal; "wGA" denotes part of the wild-type *gag* gene which contains sequences believed to be necessary for encapsidation; "X" indicates that the ATG codon of the partial *gag* gene sequence is mutated so that translation of this gene does not occur; "CMV" denotes the cytomegalovirus promoter and luciferase is used as a reporter gene. Luciferase can be replaced with any gene of interest. Another HIV LTR is present at the 3' end of transfer construct 1. Replacement of this LTR in constructs such as the transfer construct 1, 2, or 3 with a promoter-enhancer deleted HIV LTR leads to inactivation of LTR after integration. Transfer construct 2 is similar to transfer construct 1, the difference being that a mutated part of the *gag* gene (denoted "mGa") is used instead of the wild-type part of the *gag* gene. Transfer construct 3 (pm2BCwCNluci) has different mutations at the 5' splice site and has an intact ATG codon so that translation of part of the mutated *gag* gene occurs. Transfer construct 3 also has a 5' CMV promoter instead of a 5' LTR promoter. This construct is expressed independent of the presence of HIV Tat protein. The transfer constructs expressed from the LTR promoter are partially dependent on Tat protein. In 293 cells significant expression can be achieved in the absence of Tat. See, e.g., Valentin et al., Proc. Natl Acad. Sci. U S A. 95:8886-91 (1988).

#### EXAMPLE 5

##### Generation of Packaging Construct pCMVgagpol BNkan

Figure 6 shows a schematic map of the packaging construct pCMVgagpolBNKan. The nucleotide numbering is that of the HXB2R sequence



- 31 -

(Genbank accession number K03455 and M38432), where +1 is the start of transcription.

The sequence in HIV-1 *gag/pol* region was mutated in order to eliminate all the INS. The fragment from the beginning of *gag* to BsrGI site in *pol*, and the fragment KE [KpnI(3700)-EcoRI(4194)] were previously mutated described in Schneider et al., J Virol. 71: 4892-4903 (1997) and in U.S. Patent Nos. 5,972,596 and 5,965,726.

To generate pCMVgagpolBNkan, three fragments within HIV-1 *pol* region were mutated. They are fragment BP [BsrGI(2207)-PflMI(3032)], fragment PK [PflMI(3032)-KpnI(3700)] and fragment EN [EcoRI(4194)-NdeI(4668)]. Mutagenesis was performed using a modified version of the method described by Hø et al., Gene 77: 51-59 (1989) and DNA shuffling (Zhao and Arnold, Nucl. Acid Res. 25(6), 1307-1308 (1997). Sixteen oligonucleotides extending over the complete sequence of the three fragments were designed. Six oligos corresponded to fragment BP, six to fragment PK, and four to fragment EN (the oligonucleotides ranged from 130 to 195 bases in length; adjacent oligos overlapped by twenty nucleotides). Each fragment was assembled in two steps:

- 1) PCR; the reaction was carried out in standard *pfu* buffer with 10 pmol of each purified big oligo, 0.2 mM of each dNTPs and 2.5 u *pfu* DNA polymerase enzyme (Stratagene) in a 50 µl final volume. The PCR program was: 3 min 96°C followed by 50 cycles of 1 min 94°C, 1 min 55°C, and 1 min + 5 s/cycle 72°C, ended by 7 min at 72°C. After PCR, the big oligonucleotides were removed from the assembled mutated fragment.

- 2) The second step was to specifically amplify the assembled products with 30 mer primers located at the 5' and 3' end of each mutated fragment. One microliter of the assembled PCR product was used as template in a 25-cycle PCR reaction with 50 pmol of each primer, 1 x *pfu* buffer, 0.2 mM of each dNTP and 2.5 u *pfu* DNA polymerase in a 50 µl final volume. The PCR program was: 3 min 96°C, 10 cycles of 30 s 94°C, 30 s 55°C, 45 s 72°C, followed by another 14 cycles of 30 s 94°C, 30 s 55°C, 45 s + 20 s/cycle 72°C, and finally 7 min 72°C. This program gave a single PCR product of the correct size. The amplified BP, PK

- 32 -

and EN fragments were individually cloned into PCR-script™ vector using PCR-script™ Amp SK(+) Cloning Kit (Stratagene). Clones were randomly selected and sequenced. The correct BP, PK and EN fragments together with fragment KE previously mutated by Schneider et al. were ligated between BsrGI and KpnI site of p55AM1-R5 (which was previously described in Schneider et al., J. Virol. 71: 4892-4903 (1997)) to produce a completely mutated *gag/pol* ORF. The new plasmid containing the completely mutated *gag/pol* was named pLTRgagpolBN. BN stands for the modification of the fragment between BsrGI and NdeI. The mutated *gag/pol* was then cloned into a CMVkan vector containing the cytomegalovirus major late promoter (GenBank accession no. X17403) and the kanamycin resistance gene, resulting in pCMVgagpolBNkan. The plasmid backbone comes from pVR1332 provided by Vical Inc., and described in Hartikka et al., Hum Gene Ther. 7:1205-17 (1996).

It is understood that different plasmid backbones can be used, e.g., to provide good expression *in vivo*, in the case of DNA injection, for example.

#### EXAMPLE 6

##### Construction of Transfer Vectors pmBCwCNluci and pmBCmCNluci

The HIV-1 sequence BC, between BssHII (257) and ClaI (376), contains the major splice donor site and the encapsidation signal. Six oligos (33 to 46 bases) were designed to introduce mutations on the splice donor site and the AUG start codon of *gag*. The BC fragment was assembled, amplified and sequenced as described in the section concerning the construction of pCMVgagpolBN.

The mutated BC fragment and a fragment of wild type *gag* between ClaI (376) and Nsi (793) were placed between the BssHII and Nsi sites of p55RRE (Schneider et al., J. Virol. 71:4892-4903 (1997)) to generate pmBCwCN. In parallel, the fragment between ClaI (376) and NsiI sites of mutated *gag* from p55BM1-10SD+ was used to generate pmBCmCN. (p55BM1-10SD+ is similar to p55BM1-10, which is described in Schneider et al. (1997), but contains in addition the intact splice donor and encapsidation site upstream of *gag*). The region between

- 33 -

NsiI and XhoI containing 3' part of *gag* and RRE in pmBCwCN and pmBCmCN was replaced by a ClaI-XhoI fragment containing CMV promoter and luciferase gene from pHR'-CMVluci (vector from D. Trono) to generate pmBCwCNluci and pmBCmCNluci (which are shown as transfer constructs 1 and 2 in Figure 5, and schematically depicted in Figures 7 and 8, respectively). The sequences of these plasmids are shown in Figures 10 and 11, respectively. Different versions of these plasmids have also been created, by standard procedures, with variations in the region of the encapsidation site, the first splice donor site, and the initiator *gag* AUG. For example, the transfer construct pm2BcwCNluci (which is shown as transfer construct 3 in Fig. 5) has different mutations in the 5' splice site region and has an intact ATG. A comparison of the sequences in the BssHII-Cla I region of transfer constructs 1 and 2 (mBCwCN frag), transfer construct 3 (m2BCwCN frag), HXB2 and NL43 is shown in Fig. 12.

#### EXAMPLE 7

##### Preparation of Viral Particles

Lentiviral particles were generated by transient cotransfection of 293 human kidney cells with a combination of three plasmids: pCMVgagpolBNkan, pmBCwCNluci or pmBCmCNluci (transfer vector) and pHCMV-G (Yee et al., Proc. Natl. Acad. Sci., USA, 91:9564-9568 (1994) a plasmid coding for the envelope VSV-G (glycoprotein of vesicular stomatitis virus).

The day before the transfection, 293 cells were plated at a density of  $10^6$  cells/plate on a 60 mm plate. Plasmid DNA was transfected by the Ca-phosphate precipitation method in the following proportions: 3  $\mu$ g packaging construct, 6  $\mu$ g transfer construct and 100 ng VSV-G encoding construct, pHCMV-G. [Note that the LTR promoter can be expressed in 293 cells in the absence of Tat with a moderate decrease in efficiency. The transfer constructs can be fully Tat independent after replacement of the LTR promoter with a CMV (see, e.g., transfer construct 3 in Fig. 5) or other promoter in such a way that the mRNA start site is at the beginning of the LTR R region.] In the present experiments for preparation of viral particles 500 ng of a Tat expression plasmid was included in the transfection.

- 34 -

Cells were washed the day after transfection and were kept in DMEM medium for another 48 hours before the supernatants were harvested. Supernatants were spun at 1,200 rpm for 7 mins to eliminate any floating cells. pCMVgagpolBNkan produces high levels of Gag protein that is efficiently released from the cells (Figure 13), and also produces high levels of functional Pol as judged by levels of reverse transcriptase activity similar to those found upon expression of complete HIV-1 (Figure 14).

Supernatants from 293 transfected cells were used to transduce several human cell lines (293, Jurkat, U937) and non-dividing human primary macrophages.

#### EXAMPLE 8

##### Cell Transduction

Transduction was performed by incubating for 3-4 hours at 37°C the target cells with 1-2 ml of supernatant containing the retroviral vectors. The amount of retroviral vector present in the supernatant was normalized by p24 content (measured by ELISA). Equal amounts of p24 gag protein were used for infection of cells. This way, differences in production of the different preparations was minimized.

The macrophages used for transduction were isolated from the peripheral blood of healthy donors by adherence to plastic. Cells were cultured in RPMI + 20% fetal calf serum (FCS) + 10% human serum (HS). After 1 week, non-adherent cells were washed off with PBS and the macrophages were kept in culture for another 1-2 weeks in the absence of human serum. The cells were washed 2-4 times with PBS before transduction.

Cells were harvested 48 hours after transduction (seven days for primary macrophages) and the transduction efficiency was determined by measuring luciferase activity in cell extracts from the cultures. The results of the transduction experiments in 293 Jurkat, U937 and primary macrophages are shown in Figure 15A-D. These results demonstrate that Rev-independent gag-HIV-1 based retroviral vectors display high transduction efficiency in (A) 293 cells, (B) human

- 35 -

lymphoid cells, (C) human myeloid cells (U937), as well as (D) non-dividing cells such as primary human macrophages.

#### EXAMPLE 9

##### Use Of Nucleic Acids of the Invention In Immunoprophylaxis Or Immunotherapy

In postnatal gene therapy, new genetic information has been introduced into tissues by indirect means such as removing target cells from the body, infecting them with viral vectors carrying the new genetic information, and then reimplanting them into the body; or by direct means such as encapsulating formulations of DNA in liposomes; entrapping DNA in proteoliposomes containing viral envelope receptor proteins; calcium phosphate co-precipitating DNA; and coupling DNA to a polylysine-glycoprotein carrier complex. In addition, in vivo infectivity of cloned viral DNA sequences after direct intrahepatic injection with or without formation of calcium phosphate coprecipitates has also been described. mRNA sequences containing elements that enhance stability have also been shown to be efficiently translated in Xenopus laevis embryos, with the use of cationic lipid vesicles. See, e.g., J.A. Wolff, et al., Science 247:1465-1468 (1990) and references cited therein.

Recently, it has also been shown that injection of pure RNA or DNA directly into skeletal muscle results in significant expression of genes within the muscle cells. J.A. Wolff, et al., Science 247:1465-1468 (1990). Forcing RNA or DNA introduced into muscle cells by other means such as by particle-acceleration (N. -S. Yang, et al. Proc. Natl. Acad. Sci. USA 87:9568-9572 (1990); S.R. Williams et al., Proc. Natl. Acad. Sci. USA 88:2726-2730 (1991)) or by viral transduction should also allow the DNA or RNA to be stably maintained and expressed. In the experiments reported in Wolff et al., RNA or DNA vectors were used to express reporter genes in mouse skeletal muscle cells, specifically cells of the quadriceps muscles. Protein expression was readily detected and no special delivery system was required for these effects. Polynucleotide expression was also obtained when the composition and volume of the injection fluid and the method of injection were modified from the described protocol. For example, reporter enzyme activity was

- 36 -

reported to have been observed with 10 to 100  $\mu$ l of hypotonic, isotonic, and hypertonic sucrose solutions, Opti-MEM, or sucrose solutions containing 2mM  $\text{CaCl}_2$  and also to have been observed when the 10- to 100-  $\mu$ l injections were performed over 20 min. with a pump instead of within 1 min.

Enzymatic activity from the protein encoded by the reporter gene was also detected in abdominal muscle injected with the RNA or DNA vectors, indicating that other muscles can take up and express polynucleotides. Low amounts of reporter enzyme were also detected in other tissues (liver, spleen, skin, lung, brain and blood) injected with the RNA and DNA vectors. Intramuscularly injected plasmid DNA has also been demonstrated to be stably expressed in non-human primate muscle. S. Jiao et al., Hum. Gene Therapy 3:21-33 (1992).

It has been proposed that the direct transfer of genes into human muscle in situ may have several potential clinical applications. Muscle is potentially a suitable tissue for the heterologous expression of a transgene that would modify disease states in which muscle is not primarily involved, in addition to those in which it is. For example, muscle tissue could be used for the heterologous expression of proteins that can immunize, be secreted in the blood, or clear a circulating toxic metabolite. The use of RNA and a tissue that can be repetitively accessed might be useful for a reversible type of gene transfer, administered much like conventional pharmaceutical treatments. See J.A. Wolff, et al., Science 247:1465-1468 (1990) and S. Jiao et al., Hum. Gene Therapy 3:21-33 (1992).

It had been proposed by J.A. Wolff et al., supra, that the intracellular expression of genes encoding antigens might provide alternative approaches to vaccine development. This hypothesis has been supported by a recent report that plasmid DNA encoding influenza A nucleoprotein injected into the quadriceps of BALB/c mice resulted in the generation of influenza A nucleoprotein-specific cytotoxic T lymphocytes (CTLs) and protection from a subsequent challenge with a heterologous strain of influenza A virus, as measured by decreased viral lung titers, inhibition of mass loss, and increased survival. J. B. Ulmer et al., Science 259:1745-1749 (1993).

- 37 -

Therefore, it appears that the direct injection of RNA or DNA vectors encoding the viral antigen can be used for endogenous expression of the antigen to generate the viral antigen for presentation to the immune system without the need for self-replicating agents or adjuvants, resulting in the generation of antigen-specific CTLs and protection from a subsequent challenge with a homologous or heterologous strain of virus.

CTLs in both mice and humans are capable of recognizing epitopes derived from conserved internal viral proteins and are thought to be important in the immune response against viruses. By recognition of epitopes from conserved viral proteins, CTLs may provide cross-strain protection. CTLs specific for conserved viral antigens can respond to different strains of virus, in contrast to antibodies, which are generally strain-specific.

Thus, direct injection of RNA or DNA encoding the viral antigen has the advantage of being without some of the limitations of direct peptide delivery or viral vectors. See J.A. Ulmer et al., *supra*, and the discussions and references therein). Furthermore, the generation of high-titer antibodies to expressed proteins after injection of DNA indicates that this may be a facile and effective means of making antibody-based vaccines targeted towards conserved or non-conserved antigens, either separately or in combination with CTL vaccines targeted towards conserved antigens. These may also be used with traditional peptide vaccines, for the generation of combination vaccines. Furthermore, because protein expression is maintained after DNA injection, the persistence of B and T cell memory may be enhanced, thereby engendering long-lived humoral and cell-mediated immunity.

1. Vectors for the immunoprophylaxis or immunotherapy against HIV-1

The mutated *gag*, *pol* or *gag/pol* sequences will be inserted in expression vectors using a strong constitutive promoter such as CMV or RSV, or an inducible promoter such as HIV-1.

The vector will be introduced into animals or humans in a pharmaceutically acceptable carrier using one of several techniques such as injection of DNA directly into human tissues; electroporation or transfection of the

- 38 -

DNA into primary human cells in culture (*ex vivo*), selection of cells for desired properties and reintroduction of such cells into the body, (said selection can be for the successful homologous recombination of the incoming DNA to an appropriate preselected genomic region); generation of infectious particles containing the *gag* gene, infection of cells *ex vivo* and reintroduction of such cells into the body; or direct infection by said particles in vivo.

Substantial levels of protein will be produced leading to an efficient stimulation of the immune system.

In another embodiment of the invention, the described constructs will be modified to express mutated Gag proteins that are unable to participate in virus particle formation. It is expected that such Gag proteins will stimulate the immune system to the same extent as the wild-type Gag protein, but be unable to contribute to increased HIV-1 production. This modification should result in safer vectors for immunotherapy and immunoprophylaxis.

#### EXAMPLE 10

##### Inhibition of HIV-1 Expression Using Transdominant (TD)-TD-Gag-TD Rev or Td Gag-Pro-TD Rev Genes

Direct injection of DNA or use of vectors other than retroviral vectors will allow the constitutive high level of trans-dominant Gag (TDgag) in cells. In addition, the approach taken by B.K. Felber et al., *Science* 239:184-187 (1988) will allow the generation of retroviral vectors, e.g. mouse-derived retroviral vectors, encoding HIV-1 TDgag, which will not interfere with the infection of human cells by the retroviral vectors. In the approach of Felber, et al., *supra*, it was shown that fragments of the HIV-1 LTR containing the promoter and part of the polyA signal can be incorporated without detrimental effects within mouse retroviral vectors and remain transcriptionally silent. The presence of Tat protein stimulated transcription from the HIV-1 LTR and resulted in the high level expression of genes linked to the HIV-1 LTR.

The generation of hybrid TDgag-TDRev or TDgag-pro-TDRev genes and the introduction of expression vectors in human cells will allow the efficient production of two proteins that will inhibit HIV-1 expression. The incorporation of



- 39 -

two TD proteins in the same vector is expected to amplify the effects of each one on viral replication. The use of the HIV-1 promoter in a matter similar to one described in B.K. Felber, et al., supra, will allow high level Gag and Rev expression in infected cells. In the absence of infection, expression will be substantially lower. Alternatively, the use of other strong promoters will allow the constitutive expression of such proteins. This approach could be highly beneficial, because of the production of a highly immunogenic gag, which is not able to participate in the production of infectious virus, but which, in fact, antagonizes such production. This can be used as an efficient immunoprophylactic or immunotherapeutic approach against AIDS.

Examples of trans-dominant mutants are described in Trono et al., Cell 59:112-120 (1989).

1. Generation of constructs encoding transdominant Gag mutant proteins

Gag mutant proteins that can act as trans-dominant mutants, as described, for example, in Trono et al., supra, will be generated by modifying vector p37M1-10D or p55M1-13P0 to produce transdominant Gag proteins at high constitutive levels.

The transdominant Gag protein will stimulate the immune system and will inhibit the production of infectious virus, but will not contribute to the production of infectious virus.

The added safety of this approach makes it more acceptable for human application.

**VII. REFERENCES**

U.S. Patent No. 5,972,596 issued October 26, 1999 (Pavlakakis and Felber)

U.S. Patent No. 5,965,726 issued October 12, 1999 (Pavlakakis and Felber)

WO 98/17816 Lentiviral Vectors (Kingsman & Kingsman) (Oxford Biomedica Ltd)

WO 98/34640 (Shiver, J.W., Davies, M-E M., Freed, D.C., Liu, M.A. and Perry, H.C. - Merck & Co., Inc.)

WO 98/46083 Use of Lentiviral Vectors for Antigen Presentation in Dendritic Cells (Wong-Staal, Li; Kan-Mitchell) (Univ. of Cal.)

WO 99/04026 Lentiviral Vectors (Chen, Gasmi, Yee and Jolly) (Chiron)

WO 99/15641 Non-Primate Lentiviral Vectors and Packaging Systems (Poeschia, Looney and Wong-Staal) (Univ. of Cal.)

WO 99/30742 Therapeutic Use of Lentiviral Vectors (Naldini and Song)

WO 99/51754 Infectious Pseudotyped Lentiviral Vectors Lacking Matrix Protein and Uses Thereof (Goettlinger, Reil and Bukovsky) (Dana Farber Cancer Inst Inc)

- 41 -

PCT/US99/11082 Post-Transcriptional Regulatory Elements and Uses Thereof (Pavlakakis and Nappi), filed May 22, 1999, published as WO 99/61596 on December 2, 1999

Akkina, R.K., Walton, R.W., Chen, M.L., Li, Q-X, Planelles, V and Chen, I.S.Y., "High-efficiency gene transfer into CD34<sup>+</sup> cells with a human immunodeficiency virus type 1-based retroviral vector pseudotyped with vesicular stomatitis virus envelope glycoprotein G," *J. Virol.* 70:2581-2585 (1996)

Amado, R.G. & Chen, I.S.Y., "Lentiviral vectors—the promise of gene therapy within reach?," *Science* 285:674-676 (July 1999)

Donahue, R.E., An, D.S., Wersto, R.P., Agricola, B.A., Metzger, M.E. and Chen, I.S.Y., "Transplantation of immunoselected CD34<sup>+</sup> cells transduced with a EGFP-expressing lentiviral vector in non-human primates," *Blood* 92(suppl. 1):383b, Abstract #4648.5 (1998)

Fox, J.L., "Researchers wary of fear-based ban on lentivirus gene therapy," *Nature Biotechnology* 16:407-408 (1998)

Goldman, M.J., Lee, P.S., Yang, J.S. & Wilson, J.M., "Lentiviral vectors for gene therapy of cystic fibrosis," *Hum Gene Ther.* 8, 2261-2268 (1997)

Hartikka J, Sawdey M, Cornefert-Jensen F, Margalith M, Barnhart K, Nolasco M, Vahlsing HL, Meek J, Marquet M, Hobart P, Norman J, and Manthorpe M., "An improved plasmid DNA expression vector for direct injection into skeletal muscle," *Hum Gene Ther.* 7:1205-17 (1996)

Kafri, T., Blomer, U., Peterson, D.A., Gage, F.H. & Verma, I.M., "Sustained expression of genes delivered directly into liver and muscle by lentiviral vectors," *Nat Genet.* 17, 314-317 (1997)

- 42 -

Kafri, T., van Praag, H., Ouyang, L., Gage, F.G. and Verma, I.M., "A packaging cell line for lentivirus vectors," *J. Virol.* 73:576-584 (1999)

Kim, V.N., Mitrophanous, K., Kingsman, S.M., and Kingsman, A.J., "Minimal Requirement for a Lentivirus Vector Based on Human Immunodeficiency Virus Type 1", *J. Virol.* 72:811-816 (1998)

Klimatcheva, E., Rosenblatt, J.D. and Planelles, V., "Lentiviral vectors and gene therapy," *Frontiers in Bioscience* 4:d481-496 (June 1999)

Miyoshi, H., Takahashi, M., Gage, F.H. & Verma, I.M., "Stable and efficient gene transfer into the retina using an HIV-based lentiviral vector," *Proc Natl Acad Sci USA.* 94: 10319-10323 (1997)

Miyoshi, H., Blomer, U., Takahashi, M., Gage, F.H., and Verma, I.M., "Development of self-inactivating lentivirus vector," *J Virol.* 72:8150-8157 (1998)

Miyoshi, H., Smith, K.A., Mosier, D.E., Verma, I.M. and Torbett, B.E., "Transduction of human CD34<sup>+</sup> cells that mediate long-term engraftment of NOD/SCID mice by HIV vectors," *Science* 283:682-686 (1999)

Naldini, L., Blomer, U., Gallay, P., Ory, D., Mulligan, R., Gage, F.H., Verma, I.M. & Trono, D., "In vivo gene delivery and stable transduction of nondividing cells by a lentiviral vector," *Science.* 272, 263-267 (1996)

Naviaux, R.K., Costanzi, E., Haas, M. and Verma, I., "The pCL vector system: rapid production of helper-free, high-titer, recombinant retroviruses," *J. Virol.* 70:5701-5705 (1996)

Pavlakakis, G.N., Schneider, R.; Song, S., Nasioulas, G., Zolotukhin, A., Felber, B.K., Trauger, R., Cox, J., and Manthorpe, M., "Use of simple Rev-

- 43 -

independent HIV-1 gag expression vectors in gene therapy and gene vaccine applications," *Natl Conf Hum Retroviruses Relat Infect (2nd)*, Jan 29-Feb 2 (1995); 91.

Poeschla, E.M., Wong-Staal, F. & Looney, D.J., "Efficient transduction of nondividing human cells by feline immunodeficiency virus lentiviral vectors," *Nature Med.* 4:354-357 (1998)

Qiu, J. T., R. Song, M. Dettenhofer, C. Tian, T. August, B. K. Felber, G. N. Pavlakis and X. F. Yu, "Evaluation of novel human immunodeficiency virus type 1 Gag DNA vaccines for protein expression in mammalian cells and induction of immune responses," *J Virol.* 73: 9145-52 (Nov. 1999)

Reynolds, P.N. and Curiel, D.T., "Viral vectors show promise in Colorado," *Nature Biotechnology* 16:422-423 (1998)

Schneider, R., Campbell, M., Nasioulas, G., Felber, B.K., and Pavlakis, G.N., Inactivation of the human immunodeficiency virus type 1 inhibitory elements allows Rev-independent expression of Gag and Gag/protease and particle formation, "*J. Virol.* 71:4892-4903 (1997)

Schwartz, S., M. Campbell, G. Nasioulas, J. Harrison, B. K. Felber and G. N. Pavlakis, "Mutational inactivation of an inhibitory sequence in human immunodeficiency virus type-1 results in Rev-independent gag expression," *J. Virol.* 66:7176-7182 (1992)

Shiver, J.W., Yasutomi, Y., Free, D.C., Davies, M.-E., Perry, H.C., Pavlakis, G.N., Letvin, N.L., and Liu, M.A., "DNA Vaccine-Mediated Cellular Immunity Against HIV-1 gag and env", presented at the Conference on Advances in AIDS Vaccine Development: 8<sup>th</sup> Annual Meeting of the National Cooperative Vaccine Development Groups for AIDS (NCVDGs) from February 11-15, 1996.

- 44 -

Soneoka, Y., Cannon, P.M., Ransdale, E.E., Griffiths, J.C., Romano, G., Kingsman, S.M. and Kingsman, A.J., "A transient three-plasmid expression system for the production of high titer retroviral vectors," *Nuc. Acids Res.* 23:628-633 (1995).

Srinivasakumar, N., Chazal, N., Helga-Maria, C., Prasad, S., Hammar skjöld, M.-L., and Rekosh, D., "The Effect of Viral Regulatory Protein Expression on Gene Delivery by Human Immunodeficiency Virus Type 1 Vectors Produced in Stable Packaging Cell Lines," *J. Virol.*, 71:5841-5848 (1997)

Sutton, R.E., Wu, H.T., Rigg, R., Bohnlein, E. & Brown, P.O., "Human immunodeficiency virus type 1 vectors efficiently transduce human hematopoietic stem cells," *J. Virol.* 72, 5781-5788 (1998)

Tabernero, C., A. S. Zolotukhin, J. Bear, R. Schneider, G. Karsenty and B. K. Felber, "Identification of an RNA sequence within an intracisternal-A particle element able to replace Rev-mediated posttranscriptional regulation of human immunodeficiency virus type 1," *J Virol.* 71:95-101 (1997). . (see also my email message)

Takahashi, M.; Miyoshi, H.; Verma, I.M.; Gage, F.H., "Rescue from photoreceptor degeneration in the rd mouse by human immunodeficiency virus vector-mediated gene transfer," *J. Virol.* 73: 7812-7816 (Sept. 1999)

Uchida, N., Sutton, R.E., Frier, A.M., He, D., Reitsma, M.J., Chang, W.C., Veres, G., Scollay, R. & Weissman, I.L., "HIV, but not murine leukemia virus, vectors mediate high efficiency gene transfer into freshly isolated G0/G1 human hematopoietic stem cells," *Proc. Natl Acad. Sci. U.S.A.* 95, 11939-11944 (1998)

Valentin, A., W. Lu, M. Rosati, R. Schneider, J. Albert, A. Karlsson and G. N. Pavlakis. "Dual effect of interleukin 4 on HIV-1 expression: Implications

- 45 -

for viral phenotypic switch and disease progression," *Proc. Natl Acad. Sci. U S A.* 95: 8886-91 (1998)

White, S.M., Renda, M, Nam, N-Y, Klimatcheva, E., Hu, Y, Fisk, J, Halterman, M, Rimel, B.J., Federoff, H, Pandya, S., Rosenblatt, J.D. and Planelles, V, "Lentivirus vectors using human and simian immunodeficiency virus elements," *J. Virol.* 73:2832-2840 (Apr. 1999)

Wolff, J.A. and Trubetskoy, V.S., "The Cambrian period of nonviral gene delivery," *Nature Biotechnology* 16:421-422 (1998)

Zolotukhin, J., Valentin, A., Pavlakis, G. N. and Felber, B. K. "Continuous propagation of RRE(-)and Rev(-)RRE(-) human immunodeficiency virus type 1 molecular clones containing a *cis*-acting element of Simian retrovirus type 1 in human peripheral blood lymphocytes," *J. Virol.* 68:7944-7952 (1994)

Zufferey, R., Nagy, D., Mandel, R.J., Naldini, L. and Trono, D., "Multiply Attenuated Lentiviral Vector Achieves Efficient Gene-Delivery In Vivo", *Nature Biotechnology* 15:871-875 (1997)

Zufferey, R., Dull, T., Mandel, R.J., Bukovsky, A., Quiroz, D., Naldini, L. & Trono, D., "Self-inactivating lentivirus vector for safe and efficient in vivo gene delivery," *J. Virol.* 72:9873-9880 (1998)

- 46 -

Those skilled in the art will recognize that any gene encoding a mRNA containing an inhibitory/instability sequence or sequences can be modified in accordance with the exemplified methods of this invention or their functional equivalents.

Modifications of the above described modes for carrying out the invention that are obvious to those of skill in the fields of genetic engineering, virology, immunology, medicine, and related fields are intended to be within the scope of the following claims.

Every reference cited hereinbefore throughout the application is hereby incorporated by reference in its entirety.



- 47 -

WHAT IS CLAIMED IS:

1. A nucleic acid construct comprising a HIV-1 *gag/pol* gene having the coding sequence of the *gag/pol* gene set forth in Figure 1.
2. A nucleic acid construct comprising a HIV-1 *pol* gene having the coding sequence of the *pol* gene set forth in Figure 2.
3. A nucleic acid construct comprising a SIV-1 *gag* gene having the coding sequence of the *gag* gene set forth in Figure 3.
4. A nucleic acid construct comprising an HIV or SIV 5' LTR, a packaging signal, a *gag/pol* gene comprising the sequence set forth in Figure 1, a 5' splice site, a 3' splice site, an *env* gene, a *tat* gene, a functional RNA transport element and a 3' HIV or SIV LTR, said nucleic acid construct being able to produce functional Gag, Pol and Env virion components.
5. A vector comprising the nucleic acid construct of Claim 1, 2, 3 or 4.
6. A transformed host cell comprising the nucleic acid construct of Claim 1, 2, 3 or 4.
7. A transformed host cell of Claim 6 wherein said cell is a eukaryote.
8. The host cell of Claim 7 wherein said cell is a human cell.
9. A transformed host cell of Claim 6 wherein said cell is a prokaryote.
10. The host cell of Claim 9 wherein said cell is *E. coli*.
11. A pharmaceutical composition comprising the nucleic acid construct of Claim 1, 2, 3 or 4 and a pharmaceutically acceptable carrier.
12. A method for inducing antibodies in a mammal comprising administering to a mammal a composition of claim 11, wherein said nucleic acid construct is present in an amount which is effective to induce said antibodies in said mammal.
13. A method for inducing cytotoxic T lymphocytes in a mammal comprising administering to a mammal a composition of claim 11, wherein said

- 48 -

nucleic acid construct is present in an amount which is effective to induce cytotoxic T lymphocytes in said mammal.

14. A vaccine composition for inducing immunity in a mammal against HIV infection comprising a pharmaceutically acceptable carrier and further comprising a therapeutically effective amount of a nucleic acid construct of Claim 1 capable of producing HIV Gag and Pol proteins in the absence of HIV Rev regulatory protein in a cell in vivo.

15. A vaccine composition for inducing immunity in a mammal against HIV infection comprising a pharmaceutically acceptable carrier and further comprising a therapeutically effective amount of a nucleic acid construct of Claim 2 capable of producing HIV Pol protein in the absence of HIV Rev regulatory protein in a cell in vivo.

16. A vaccine composition according to claim 14 wherein said mammal is a human.

17. A vaccine composition according to claim 15 wherein said mammal is a human.

18. A method for inducing immunity against HIV infection in a mammal which comprises administering to a mammal a therapeutically effective amount of a vaccine composition according to claim 14.

19. A method for inducing immunity against HIV infection in a mammal which comprises administering to a mammal a therapeutically effective amount of a vaccine composition according to claim 15.

20. A method according to claim 18 wherein said mammal is a human.

21. A method according to claim 19 wherein said mammal is a human.

22. A lentiviral expression system comprising the following:

- (a) a packaging vector comprising a HIV-1 *gag/pol* gene having the nucleotide sequence set forth in Figure 1;
- (b) a transfer vector; and
- (c) an envelope encoding vector.

- 49 -

23. A transformed host cell comprising the lentiviral expression system of Claim 22.

24. A transformed host cell of Claim 23 wherein said cell is a eukaryote.

25. The host cell of Claim 24 wherein said cell is a human cell.

26. A process for making a lentiviral particle comprising expressing HIV Gag and HIV Pol in a host cell from a vector comprising the nucleotide sequences encoding HIV Gag and HIV Pol set forth in Figure 1 in the presence of a gene encoding an envelope protein.

27. A lentiviral expression system which is capable of functioning in the absence of Rev, Tat, and any viral RNA transport element comprising the following:

(a) a packaging vector comprising a HIV-1 *gag/pol* gene which has been mutated to eliminate inhibitory/instability regions;

(b) a transfer vector; and

(c) an envelope encoding vector.

28. A transformed host cell comprising the lentiviral expression system of Claim 27.

29. A transformed host cell of Claim 28 wherein said cell is a eukaryote.

30. The host cell of Claim 29 wherein said cell is a human cell.

31. A process for making a lentiviral particle in the absence of Rev, Tat, or any viral RNA transport element comprising expressing HIV Gag and HIV Pol in a host cell from a HIV-1 *gag/pol* gene which has been mutated to eliminate inhibitory/instability regions and expressing an Envelope protein from a envelope encoding gene whose expression is independent Rev, Tat, or any viral RNA transport element.

32. A nucleic acid construct comprising a SIV-1 *env* gene having the coding sequence of the *env* gene set forth in Figure 16.

33. A vector comprising the nucleic acid construct of claim 32.

34. A transformed host cell comprising the nucleic acid construct of claim 32.

- 50 -

35. A pharmaceutical composition comprising the nucleic acid construct of claim 32 and a pharmaceutically acceptable carrier.

1/43

## FIGURE 1

ATGGGTGCGAGAGCGTCAGTATTAAGCGGGGAGAATTAGATCGATGGGAAAAAAT  
TCGGTTAAGGCCAGGGGAAAGAAGAAGTACAAGCTAAAGCACATCGTATGGGCAA  
GCAGGGAGCTAGAACGATTCGCAGTTAATCCTGGCCTGTTAGAAACATCAGAAGGC  
TGTAGACAAATACTGGGACAGCTACAACCATCCCTTCAGACAGGATCAGAGGAGCT  
TCGATCACTATACAACACAGTAGCAACCCTCTATTGTGTGCACCAGCGGATCGAGA  
TCAAGGACACCAAGGAAGCTTTAGACAAGATAGAGGAAGAGCAAAACAAGTCCAAG  
AAGAAGGCCCCAGCAGGCAGCAGCTGACACAGGACACAGCAATCAGGTCAGCCAAAA  
TTACCCTATAGTGCAGAACATCCAGGGGCAAATGGTACATCAGGCCATATCACCTA  
GAACTTTAAATGCATGGGTAAAAGTAGTAGAAGAGAAGGCTTTCAGCCCAGAAGTG  
ATACCCATGTTTTTCAGCATTATCAGAAGGAGCCACCCACAGGACCTGAACACGAT  
GTTGAACACCGTGGGGGGACATCAAGCAGCCATGCAAAATGTTAAAAGAGACCATCA  
ATGAGGAAGCTGCAGAATGGGATAGAGTGCATCCAGTGCATGCAGGGCCTATTGCA  
CCAGGCCAGATGAGAGAACCAAGGGGAAGTGACATAGCAGGAACTACTAGTACCCT  
TCAGGAACAAATAGGATGGATGACAAATAATCCACCTATCCCAGTAGGAGAGATCT  
ACAAGAGGTGGATAATCCTGGGATTGAACAAGATCGTGAGGATGTATAGCCCTACC  
AGCATTCTGGACATAAGACAAGGACCAAAGGAACCCTTTAGAGACTATGTAGACCG  
GTTCTATAAACTCTAAGAGCTGAGCAAGCTTCACAGGAGGTAAAAAATTGGATGA  
CAGAAACCTTGTTGGTCCAAATGCGAACCAGATTGTAAGACCATCCTGAAGGCT  
CTCGGCCCAGCGGCTACACTAGAAGAAATGATGACAGCATGTCAGGGAGTAGGAGG  
ACCCGGCCATAAGGCAAGAGTTTTGGCCGAGGCGATGAGCCAGGTGACGAACTCGG  
CGACCATAATGATGCAGAGAGGCAACTTCCGGAACCAGCGGAAGATCGTCAAGTGC  
TTCAATTGTGGCAAAGAAGGGCACACCGCCAGGAACTGCCGGGGCCCCCGGAAGAA  
GGGCTGTTGGAAATGTGGAAAGGAAGGACACCAAATGAAAGATTGTACTGAGAGAC

2/43

AGGCTAATTTTTTAGGGAAGATCTGGCCTTCCTACAAGGGAAGGCCAGGGAATTTT  
CTTCAGAGCAGACCAGAGCCAACAGCCCCACCAGAAGAGAGCTTCAGGTCTGGGGT  
AGAGACAACAACCTCCCCCTCAGAAGCAGGAGCCGATAGACAAGGAACTGTATCCTT  
TAACTTCCCTCAGATCACTCTTTGGCAACGACCCCTCGTCACAGTAAGGATCGGGG  
GGCAACTCAAGGAAGCGCTGCTCGATACAGGAGCAGATGATACAGTATTAGAAGAA  
ATGAGTTTGCCAGGAAGATGGAAACCAAAAATGATAGGGGGGATCGGGGGCTTCAT  
CAAGGTGAGGCAGTACGACCAGATACTCATAGAAATCTGTGGACATAAAGCTATAG  
GTACAGTATTAGTAGGACCTACACCTGTCAACATAATTGGAAGAAATCTGTTGACC  
CAGATCGGCTGCACCTTGAACCTCCCCATCAGCCCTATTGAGACGGTGCCCGTGAA  
GTTGAAGCCGGGGATGGACGGCCCCAAGGTCAAGCAATGGCCATTGACGAAAGAGA  
AGATCAAGGCCTTAGTCGAAATCTGTACAGAGATGGAGAAGGAAGGGAAGATCAGC  
AAGATCGGGCCTGAGAACCCCTACAACACTCCAGTCTTCGCAATCAAGAAGAAGGA  
CAGTACCAAGTGGAGAAAGCTGGTGGACTTCAGAGAGCTGAACAAGAGAACTCAGG  
ACTTCTGGGAAGTTCAGCTGGGCATCCACATCCCGCTGGGTTGAAGAAGAAGAAG  
TCAGTGACAGTGCTGGATGTGGGTGATGCCTACTTCTCCGTTCCCTTGGACGAGGA  
CTTCAGGAAGTACACTGCCTTCACGATACCTAGCATCAACAACGAGACACCAGGCA  
TCCGCTACCAGTACAACGTGCTGCCACAGGGATGGAAGGGATCACCAGCCATCTTT  
CAAAGCAGCATGACCAAGATCCTGGAGCCCTTCGCAAGCAAAACCCAGACATCGT  
GATCTATCAGTACATGGACGACCTCTACGTAGGAAGTGACCTGGAGATCGGGCAGC  
ACAGGACCAAGATCGAGGAGCTGAGACAGCATCTGTTGAGGTGGGGACTGACCACA  
CCAGACAAGAAGCACCAGAAGGAACCTCCCTTCCTGTGGATGGGCTACGAACTGCA  
TCCTGACAAGTGGACAGTGCAGCCCATCGTGCTGCCTGAGAAGGACAGCTGGACTG  
TGAACGACATACAGAAGCTCGTGGGCAAGTTGAACTGGGCAAGCCAGATCTACCCA  
GGCATCAAAGTTAGGCAGCTGTGCAAGCTGCTTCGAGGAACCAAGGCACTGACAGA

3/43

AGTGATCCCACTGACAGAGGAAGCAGAGCTAGA ACTGGCAGAGAACCGAGAGATCC  
TGAAGGAGCCAGTACATGGAGTGTACTACGACCCAAGCAAGGACCTGATCGCAGAG  
ATCCAGAAGCAGGGGCAAGGCCAATGGACCTACCAAATCTACCAGGAGCCCTTCAA  
GAACCTGAAGACAGGCAAGTACGCAAGGATGAGGGGTGCCCACACCAACGATGTGA  
AGCAGCTGACAGAGGCAGTGCAGAAGATCACCACAGAGAGCATCGTGATCTGGGGC  
AAGACTCCCAAGTTCAAGCTGCCCCATACAGAAGGAGACATGGGAGACATGGTGGAC  
CGAGTACTGGCAAGCCACCTGGATCCCTGAGTGGGAGTTCGTGAACACCCCTCCCT  
TGGTGAACTGTGGTATCAGCTGGAGAAGGAACCCATCGTGGGAGCAGAGACCTTC  
TACGTGGATGGGGCAGCCAACAGGGAGACCAAGCTGGGCAAGGCAGGCTACGTGAC  
CAACCGAGGACGACAGAAAGTGGTGACCCTGACTGACACCACCAACCAGAAGACTG  
AGCTGCAAGCCATCTACCTAGCTCTGCAAGACAGCGGACTGGAAGTGAACATCGTG  
ACAGACTCACAGTACGCACTGGGCATCATCCAAGCACAACCAGACCAATCCGAGTC  
AGAGCTGGTGAACCAGATCATCGAGCAGCTGATCAAGAAGGAGAAAGTGACCTGG  
CATGGGTACCAGCACACAAAGGAATTGGAGGAAATGAACAAGTAGATAAATTAGTC  
AGTGCTGGGATCCGGAAGGTGCTGTTCTGGACGGGATCGATAAGGCCCAAGATGA  
ACATGAGAAGTACCACTCCA ACTGGCGCGCTATGGCCAGCGACTTCAACCTGCCAC  
CTGTAGTAGCAAAAGAAATAGTAGCCAGCTGTGATAAATGTCAGCTAAAAGGAGAA  
GCCATGCATGGACAAGTAGACTGTAGTCCAGGAATATGGCAGCTGGACTGCACGCA  
CCTGGAGGGGAAGGTGATCCTGGTAGCAGTTCATGTAGCCAGTGGATATATAGAAG  
CAGAAGTTATCCCTGCTGAACTGGGCAGGAAACAGCATATTTTCTTTTAAATTA  
GCAGGAAGATGGCCAGTAAAACAATACACACGGACAACGGAAGCAACTTCACTGG  
TGCTACGGTTAAGGCCGCCTGTTGGTGGGCGGGAATCAAGCAGGAATTTGGAATTC  
CCTACAATCCCCAATCGCAAGGAGTCGTGGAGAGCATGAACAAGGAGCTGAAGAAG  
ATCATCGGACAAGTGAGGGATCAGGCTGAGCACCTGAAGACAGCAGTGCAGATGGC

4/43

AGTGTTTCATCCACAACCTTCAAAAGAAAAGGGGGGATTGGGGGGTACAGTGCAGGGG  
AAAGGATCGTGGACATCATCGCCACCGACATCCAAACCAAGGAGCTGCAGAAGCAG  
ATCACCAAGATCCAGAACTTCCGGGTGTACTACCGCGACAGCCGCAACCCACTGTG  
GAAGGGACCAGCAAAGCTCCTCTGGAAGGGAGAGGGGGCAGTGGTGATCCAGGACA  
ACAGTGACATCAAAGTGGTGCCAAGGCGCAAGGCCAAGATCATCCGCGACTATGGA  
AAACAGATGGCAGGTGATGATTGTGTGGCAAGTAGACAGGATGAGGATTAGAACCT  
GGAAGAGCCTGGTGAAGCACCATATG (SEQUENCE ID NO:1)



5/43

FIGURE 2

```

>wildtype      TGTACAGAGA TGGAAAAGGA AGGGAAAATT TCAAAAATTG
>mutated       TGTACAGAGA TGGAGAAGGA AGGGAAGATC AGCAAGATCG
#1             .....
                *      * * *** * *

>wildtype      GGCCTGAAAA TCCATACAAT ACTCCAGTAT TTGCCATAAA
>mutated       GGCCTGAGAA CCCCTACAAC ACTCCAGTCT TCGCAATCAA
#41            .....
                * * * * *

>wildtype      GAAAAAAGAC AGTACTAAAT GGAGAAAATT AGTAGATTTC
>mutated       GAAGAAGGAC AGTACCAAGT GGAGAAAGCT GGTGGACTTC
#81            .....
                * * * * *

>wildtype      AGAGAACTTA ATAAGAGAAC TCAAGACTTC TGGGAAGTTC
>mutated       AGAGAGCTGA ACAAGAGAAC TCAGGACTTC TGGGAAGTTC
#121           .....
                * * * *

>wildtype      AATTAGGAAT ACCACATCCC GCAGGGTTAA AAAAGAAAAA
>mutated       AGCTGGGCAT CCCACATCCC GCTGGGTGTA AGAAGAAGAA
#161           .....
                ** * * * *

>wildtype      ATCAGTAACA GTACTGGATG TGGGTGATGC ATATTTTCA
>mutated       GTCAGTGACA GTGCTGGATG TGGGTGATGC CTACTTCTCC
#201           .....
                * * * *

>wildtype      GTTCCCTTAG ATGAAGACTT CAGGAAATAT ACTGCATTTA
>mutated       GTTCCCTTGG ACGAGGACTT CAGGAAATAC ACTGCCTTCA
#241           .....
                * * * *

>wildtype      CCATACCTAG TATAAACAAT GAGACACCAG GGATTAGATA
>mutated       CGATACCTAG CATCAACAAC GAGACACCAG GCATCCGCTA
#281           .....
                * * * *

>wildtype      TCAGTACAAT GTGCTTCCAC AGGGATGGAA AGGATCACCA
>mutated       CCAGTACAAC GTGCTGCCAC AGGGATGGAA GGGATCACCA
#321           .....
                * * *

>wildtype      GCAATATTCC AAAGTAGCAT GACAAAAATC TTAGAGCCTT
>mutated       GCCATCTTTC AAAGCAGCAT GACCAAGATC CTGGAGCCCT
#361           .....
                * * * *

>wildtype      TTAGAAAACA AAATCCAGAC ATAGTTATCT ATCAATACAT
>mutated       TCCGCAAGCA AAATCCAGAC ATCGTGATCT ATCAGTACAT
#401           .....
                ** * * *

```

6/43

```

>wildtype      GGATGATTTG TATGTAGGAT CTGACTTAGA AATAGGGCAG
>mutated       GGACGACCTC TACGTAGGAA GTGACCTGGA GATCGGGCAG
#441           .....
               *  * *  *      * *      * *      * *

>wildtype      CATAGAACAA AAATAGAGGA GCTGAGACAA CATCTGTTGA
>mutated       CACAGGACCA AGATCGAGGA GCTGAGACAG CATCTGTTGA
#481           .....
               *  *  *  *      *      *

>wildtype      GGTGGGGACT TACCACACCA GACAAAAAAC ATCAGAAAGA
>mutated       GGTGGGGACT GACCACACCA GACAAGAAGC ACCAGAAGGA
#521           .....
               *      *      *      *

>wildtype      ACCTCCATTC CTTTGGATGG GTTATGAACT CCATCCTGAT
>mutated       ACCTCCCTTC CTGTGGATGG GCTACGAACT GCATCCTGAC
#561           .....
               *      *      *      *

>wildtype      AAATGGACAG TACAGCCTAT AGTGCTGCCA GAAAAAGACA
>mutated       AAGTGGACAG TGCAGCCCAT CGTGCTGCCT GAGAAGGACA
#601           .....
               *      *      *      *

>wildtype      GCTGGACTGT CAATGACATA CAGAAGTTAG TGGGGAAATT
>mutated       GCTGGACTGT GAACGACATA CAGAAGCTCG TGGGCAAGTT
#641           .....
               *      *      *      *

>wildtype      GAATTGGGCA AGTCAGATTT ACCCAGGGAT TAAAGTAAGG
>mutated       GAACTGGGCA AGCCAGATCT ACCCAGGCAT CAAAGTTAGG
#681           .....
               *      *      *      *

>wildtype      CAATTATGTA AACTCCTTAG AGGAACCAAA GCACTAACAG
>mutated       CAGCTGTGCA AGCTGCTTCG AGGAACCAAG GCACTGACAG
#721           .....
               ** * *  * *      *      *

>wildtype      AAGTAATACC ACTAACAGAA GAAGCAGAGC TAGAACTGGC
>mutated       AAGTGATCCC ACTGACAGAG GAAGCAGAGC TAGAACTGGC
#761           .....
               *      *      *      *

>wildtype      AGAAAACAGA GAGATTCTAA AAGAACCAGT ACATGGAGTG
>mutated       AGAGAACCGA GAGATCCTGA AGGAGCCAGT ACATGGAGTG
#801           .....
               *      *      *      *

>wildtype      TATTATGACC CATCAAAGA CTTAATAGCA GAAATACAGA
>mutated       TACTACGACC CAAGCAAGGA CCTGATCGCA GAGATCCAGA
#841           .....
               *      *      *** *      * *      * *

```

7/43

```
>wildtype      AGCAGGGGCA AGGCCAATGG ACATATCAAA TTTATCAAGA
>mutated       AGCAGGGGCA AGGCCAATGG ACCTACCAAA TCTACCAGGA
#881           .....
                        * * * * *

>wildtype      GCCATTTAAA AATCTGAAAA CAGGAAAATA TGCAAGAATG
>mutated       GCCCTTCAAG AACCTGAAGA CAGGCAAGTA CGCAAGGATG
#921           .....
                        * * * * *

>wildtype      AGGGGTGCCC AACTAATGA TGTAACAA TTAACAGAGG
>mutated       AGGGGTGCCC ACACCAACGA TGTGAAGCAG CTGACAGAGG
#961           .....
                        * * * * *

>wildtype      CAGTGCAAAA AATAACCACA GAAAGCATAG TAATATGGGG
>mutated       CAGTGCAGAA GATCACCACA GAGAGCATCG TGATCTGGGG
#1001          .....
                        * * * * *

>wildtype      AAAGACTCCT AAATTTAAAC TGCCCATACA AAAGGAAACA
>mutated       CAAGACTCCC AAGTTCAAGC TGCCCATACA GAAGGAGACA
#1041          .....
                        * * * * *

>wildtype      TGGGAAACAT GGTGGACAGA GTATTGGCAA GCCACCTGGA
>mutated       TGGGAGACAT GGTGGACCGA GTACTGGCAA GCCACCTGGA
#1081          .....
                        * * * * *

>wildtype      TTCCTGAGTG GGAGTTTGT AATACCCCTC CTTTAGTGAA
>mutated       TCCCTGAGTG GGAGTTCGTG AACACCCCTC CCTTGGTGAA
#1121          .....
                        * * * * *

>wildtype      ATTATGGTAC CAGTTAGAGA AAGAACCCAT AGTAGGAGCA
>mutated       ACTGTGGTAT CAGCTGGAGA AGGAACCCAT CGTGGGAGCA
#1161          .....
                        * * * * *

>wildtype      GAAACCTTCT ATGTAGATGG GGCAGCTAAC AGGGAGACTA
>mutated       GAGACCTTCT ACGTGGATGG GGCAGCCAAC AGGGAGACCA
#1201          .....
                        * * * * *

>wildtype      AATTAGGAAA AGCAGGATAT GTTACTAATA GAGGAAGACA
>mutated       AGCTGGGCAA GGCAGGCTAC GTGACCAACC GAGGACGACA
#1241          .....
                        * * * * *

>wildtype      AAAAGTTGTC ACCCTAACTG ACACAACAAA TCAGAAGACT
>mutated       GAAAGTGGTG ACCCTGACTG ACACCACCAA CCAGAAGACT
#1281          .....
                        * * * * *
```

8/43

```

>wildtype      GAGTTACAAG CAATTTATCT AGCTTTGCAG GATTCGGGAT
>mutated       GAGCTGCAAG CCATCTACCT AGCTCTGCAA GACAGCGGAC
#1321          .....
          * * * * *

>wildtype      TAGAAGTAAA CATAGTAACA GACTCACAAT ATGCATTAGG
>mutated       TGGAAGTGAA CATCGTGACA GACTCACAGT ACGCACTGGG
#1361          .....
          * * * * *

>wildtype      AATCATTCAA GCACAACCAG ATCAAAGTGA ATCAGAGTTA
>mutated       CATCATCCAA GCACAACCAG ACCAATCCGA GTCAGAGCTG
#1401          .....
          * * * * *

>wildtype      GTCAATCAAA TAATAGAGCA GTTAATAAAA AAGGAAAAGG
>mutated       GTGAACCAGA TCATCGAGCA GCTGATCAAG AAGGAGAAAG
#1441          .....
          * * * * *

>wildtype      TCTATCTGGC ATGGGTACCA GCACACAAAG GAATTGGAGG
>mutated       TGTACCTGGC ATGGGTACCA GCACACAAAG GAATTGGAGG
#1481          .....
          * *

>wildtype      AAATGAACAA GTAGATAAAT TAGTCAGTGC TGGGAATCAGG
>mutated       AAATGAACAA GTAGATAAAT TAGTCAGTGC TGGGATCCGG
#1521          .....
          * *

>wildtype      AAAGTACTAT TTTTAGATGG AATAGATAAG GCCCAAGATG
>mutated       AAGGTGCTGT TCCTGGACGG GATCGATAAG GCCCAAGATG
#1561          .....
          * * * * *

>wildtype      AACATGAGAA ATATCACAGT AATTGGAGAG CAATGGCTAG
>mutated       AACATGAGAA GTACCACTCC AACTGGCGCG CTATGGCCAG
#1601          .....
          * * * * *

>wildtype      TGATTTTAAC CTGCCACCTG TAGTAGCAAA AGAAATAGTA
>mutated       CGACTTCAAC CTGCCACCTG TAGTAGCAAA AGAAATAGTA
#1641          .....
          * * *

>wildtype      GCCAGCTGTG ATAAATGTCA GCTAAAAGGA GAAGCCATGC
>mutated       GCCAGCTGTG ATAAATGTCA GCTAAAAGGA GAAGCCATGC
#1681          .....

>wildtype      ATGGACAAGT AGACTGTAGT CCAGGAATAT GGCAACTAGA
>mutated       ATGGACAAGT AGACTGTAGT CCAGGAATAT GGCAGCTGGA
#1721          .....
          * *

```

9/43

```
>wildtype      TTGTACACAT TTAGAAGGAA AAGTTATCCT GGTAGCAGTT
>mutated       CTGCACGCAC CTGGAGGGGA AGGTGATCCT GGTAGCAGTT
#1761          .....
          * * * * *

>wildtype      CATGTAGCCA GTGGATATAT AGAAGCAGAA GTTATTCCAG
>mutated       CATGTAGCCA GTGGATATAT AGAAGCAGAA GTTATCCCTG
#1801          .....
          * *

>wildtype      CAGAAACAGG GCAGGAAACA GCATATTTTC TTTTAAAATT
>mutated       CTGAAACTGG GCAGGAAACA GCATATTTTC TTTTAAAATT
#1841          .....
          *

>wildtype      AGCAGGAAGA TGGCCAGTAA AAACAATACA TACAGACAAT
>mutated       AGCAGGAAGA TGGCCAGTAA AAACAATACA CACGGACAAC
#1881          .....
          * * *

>wildtype      GGCAGCAATT TCACCAAGTGC TACGGTTAAG GCCGCCTGTT
>mutated       GGAAGCAACT TCACTGGTGC TACGGTTAAG GCCGCCTGTT
#1921          .....
          * * *

>wildtype      GGTGGGCGGG AATCAAGCAG GAATTTGGAA TTCCCTACAA
>mutated       GGTGGGCGGG AATCAAGCAG GAATTTGGAA TTCCCTACAA
#1961          .....

>wildtype      TCCCCAAAGT CAAGGAGTAG TAGAATCTAT GAATAAAGAA
>mutated       TCCCCAATCG CAAGGAGTCG TGGAGAGCAT GAACAAGGAG
#2001          .....
          *** * * **** * * *

>wildtype      TTAAAGAAAA TTATAGGACA GGTAAGAGAT CAGGCTGAAC
>mutated       CTGAAGAAGA TCATCGGACA AGTGAGGGAT CAGGCTGAGC
#2041          .....
          * * * * *

>wildtype      ATCTTAAGAC AGCAGTACAA ATGGCAGTAT TCATCCACAA
>mutated       ACCTGAAGAC AGCAGTGCAG ATGGCAGTGT TCATCCACAA
#2081          .....
          * * * * *

>wildtype      TTTTAAAAGA AAAGGGGGGA TTGGGGGGTA CAGTGCAGGG
>mutated       CTTCAAAAAGA AAAGGGGGGA TTGGGGGGTA CAGTGCAGGG
#2121          .....
          *

>wildtype      GAAAGAATAG TAGACATAAT AGCAACAGAC ATACAAACTA
>mutated       GAAAGGATCG TGGACATCAT CGCCACCGAC ATCCAAACCA
#2161          .....
          * * * * *
```

10/43

```
>wildtype      AAGAATTACA AAAACAAATT ACAAAAATTC AAAATTTTCG
>mutated       AGGAGCTGCA GAAGCAGATC ACCAAGATCC AGAACTTCCG
#2201          .....
                * * * * *
>wildtype      GGTTTATTAC AGGGACAGCA GAAATCCACT TTGGAAAGGA
>mutated       GGTGTACTAC CGCGACAGCC GCAACCCACT GTGGAAGGGA
#2241          .....
                * * * * *
>wildtype      CCAGCAAAGC TCCTCTGGAA AGGTGAAGGG GCAGTAGTAA
>mutated       CCAGCAAAGC TCCTCTGGAA GGGAGAGGGG GCAGTGGTGA
#2281          .....
                        * * * * *
>wildtype      TACAAGATAA TAGTGACATA AAAGTAGTGC CAAGAAGAAA
>mutated       TCCAGGACAA CAGTGACATC AAAGTGGTGC CAAGGCGCAA
#2321          .....
                * * * * *
>wildtype      AGCAAAGATC ATTAGGGATT ATGGAAAACA GATGGCAGGT
>mutated       GGCCAAGATC ATCCGCGACT ATGGAAAACA GATGGCAGGT
#2361          .....
                * * * * *
>wildtype      GATGATTGTG TGGCAAGTAG ACAGGATGAG GATTAGAACA
>mutated       GATGATTGTG TGGCAAGTAG ACAGGATGAG GATTAGAACC
#2401          .....
                                     +
>wildtype      TGGAAGAGTT TAGTAAAACA CCATATG
>mutated       TGGAAGAGCC TGGTGAAGCA CCATATG
#2441          .....
                * * * * *
```

11/43

FIGURE 3

ATGGGCGTGAGAACTCCGTCTTGTCAGGGAAGAAAGCAGATGAATTAG  
AAAAAATTAGGCTACGACCCAACGGAAAGAAAAAGTACATGTTGAAGC  
ATGTAGTATGGGCAGCAAATGAATTAGATAGATTTGGATTAGCAGAAAG  
CCTGTTGGAGAACAAAGAAGGATGTCAAAAAATACTTTCGGTCTTAGCT  
CCATTAGTGCCAACAGGCTCAGAAAATTTAAAAAGCCTTTATAATACTG  
TCTGCGTCATCTGGTGCATTACGCGAGAAGAGAAAAGTGAAACACACTGA  
GGAAGCAAAACAGATAGTGCAGAGACACCTAGTGGTGGAAACAGGAAC  
CACCGAAACCATGCCGAAGACCTCTCGACCAACAGCACCATCTAGCGGC  
AGAGGAGGAAACTACCCAGTACAGCAGATCGGTGGCAACTACGTCCAC  
CTGCCACTGTCCCCGAGAACCCTGAACGCTTGGGTCAAGCTGATCGAGG  
AGAAGAAGTTCGGAGCAGAAGTAGTGCCAGGATTCCAGGCACTGTCAG  
AAGGTTGCACCCCCTACGACATCAACCAGATGCTGAACTGCGTTGGAGA  
CCATCAGGCGGCTATGCAGATCATCCGTGACATCATCAACGAGGAGGCT  
GCAGATTGGGACTTGCAGCACCCACAACCAGCTCCACAACAAGGACAA  
CTTAGGGAGCCGTCAGGATCAGACATCGCAGGAACCACTCCTCAGTTG  
ACGAACAGATCCAGTGGATGTACCGTCAGCAGAACCCGATCCCAAGTAGG  
CAACATCTACCGTCGATGGATCCAGCTGGGTCTGCAGAAATGCGTCCGT  
ATGTACAACCCGACCAACATTCTAGATGTAAAACAAGGGCCAAAAGAG  
CCATTTTCAGAGCTATGTAGACAGGTTCTACAAAAGTTTAAGAGCAGAAC  
AGACAGATGCAGCAGTAAAGAATTGGATGACTCAAACACTGCTGATTCA  
AAATGCTAACCCAGATTGCAAGCTAGTGCTGAAGGGGCTGGGTGTGAAT  
CCCACCCTAGAAGAAATGCTGACGGCTTGTCAAGGAGTAGGGGGGGCCG  
GGACAGAAGGCTAGATTAATGGCAGAAGCCCTGAAAGAGGGCCCTCGCA  
CCAGTGCCAATCCCTTTTGCAGCAGCCCAACAGAGGGGACCAAGAAAGC  
CAATTAAGTGTTGGAATTGTGGGAAAGAGGGACACTCTGCAAGGCAATG  
CAGAGCCCCAAGAAGACAGGGATGCTGGAAATGTGGAAAAATGGACCA  
TGTTATGGCCAAATGCCCAGACAGACAGGCGGGTTTTTTAGGCCTTGGT  
CCATGGGGAAAGAAGCCCCGCAATTTCCCCATGGCTCAAGTGCATCAGG  
GGCTGATGCCAACTGCTCCCCCAGAGGACCCAGCTGTGGATCTGCTAAA  
GAACTACATGCAGTTGGGCAAGCAGCAGAGAGAAAAGCAGAGAGAAAAG  
CAGAGAGAAGCCTTACAAGGAGGTGACAGAGGATTTGCTGCACCTCAAT  
TCTCTCTTTGGAGGAGACCAGTAG

12/43

FIGURE 4

```

SIV gag  -----
#1      .....
      ATGGGCGTGAGAACTCCGTCTTGTCAGGGAAGAAAGCAG

SIV gag  -----
#41     .....
      ATGAATTAGAAAAAATTAGGCTACGACCCAACGGAAAGAA

SIV gag  -----
#81     .....
      AAAGTACATGTTGAAGCATGTAGTATGGGCAGCAAATGAA

SIV gag  -----
#121    .....
      TTAGATAGATTTGGATTAGCAGAAAGCCTGTTGGAGAACA

SIV gag  -----
#161    .....
      AAGAAGGATGTCAAAAAATACTTTCGGTCTTAGCTCCATT

SIV gag  -----
#201    .....
      AGTGCCAACAGGCTCAGAAAATTTAAAAAGCCTTTATAAT

SIV gag  -----
#241    .....
      ACTGTCTGCGTCATCTGGTGCATTCACGCAGAAGAGAAAG

SIV gag  -----
SIVgagDX.. -----
#281    .....
      TGAAACACACTGAGGAAGCAAAACAGATAGTGCAGAGACA

SIV gag  -----A--A-----T-----A--A
SIVgagDX..-----C--C-----C-----G--G
#321    .....
      CCTAGTGGTGGAAACAGGAACMACMGAAACYATGCCRAAR

SIV gag  --AAG-A-----
SIVgagDX..--CTC-C-----
#361    .....
      ACMWSTMGACCAACAGCACCATCTAGCGGCAGAGGAGGAA

```



13/43

```

SIV gag      -T-----A--A--A-----T-----
SIVgagDX..  -C-----G--G--C-----C-----
#401          .....
          AYTACCCAGTACARCARATMGGTGGTAACTAYGTCCACCT

SIV gag      ----T-AAG-----AT-A--T--C----A--AT--
SIVgagDX..  ----C-GTC-----CC-G--C--T----C--GC--
#441          .....
          GCCAYTRWSCCCGAGAACMYTRAAYGCTGGGTMAARYTG

SIV gag      --A-----A-----A--T-----
SIVgagDX..  --C-----G-----G--C-----
#481          .....
          ATMGAGGARAAGAARTTYGGAGCAGAAGTAGTGCCAGGAT

SIV gag      -T-----T-----T-----T-----
SIVgagDX..  -C-----C-----C-----C-----
#521          .....
          TYCAGGCACTGTCAGAAGGTTGCACCCCTAYGACATYAA

SIV gag      T-----T-A--T--T--G-----A-----
SIVgagDX..  C-----C-G--C--C--T-----G-----
#561          .....
          YCAGATGYTRAAYTGYGTKGGAGACCATCARGCGGCTATG

SIV gag      ----T---A-A--T--T--A-----
SIVgagDX..  ----C---C-T--C--C--C-----
#601          .....
          CAGATYATCMGWGAYATYATMAACGAGGAGGCTGCAGATT

SIV gag      -----
SIVgagDX..  -----
#641          .....
          GGGACTTGTCAGCACCCACAACCAGCTCCACAACAAGGACA

SIV gag      -----T--T-----A--T
SIVgagDX..  -----C--C-----C--C
#681          .....
          ACTTAGGGAGCCGTCAGGATCAGAYATYGCAGGAACMACY

SIV gag      AGT----A--T----A-----A-A--A-
SIVgagDX..  TCC----T--C----G-----C-T--G-
#721          .....
          WSYTCAGTWGAYGAACARATCCAGTGGATGTACMGWCARC

```

14/43

```
SIV gag      -----C--A-----T--A-GA-----
SIVgagDX..   -----G--C-----C--C-TC-----
#761          .....
              AGAACCCSATMCCAGTAGGCAACATYTACMGKMGATGGAT

SIV gag      ---A-----GT---A--A--T--CA-A-----T-----A
SIVgagDX..   ---G-----TC---G--G--C--TC-T-----C-----G
#801          .....
              CCARCTGGGKYTGCARAARTGYGTYMGWATGTAYAACCCR

SIV gag      --A-----
SIVgagDX..   --C-----
#841          .....
              ACMAACATTCTAGATGTAAAACAAGGGCCAAAAGAGCCAT

SIV gag      -----
#881          .....
              TTCAGAGCTATGTAGACAGGTTCTACAAAAGTTTAAGAGC

SIV gag      -----
#921          .....
              AGAACAGACAGATGCAGCAGTAAAGAATTGGATGACTCAA

SIV gag      -----
#961          .....
              ACACTGCTGATTCAAAATGCTAACCCAGATTGCAAGCTAG

SIV gag      -----
#1001         .....
              TGCTGAAGGGGCTGGGTGTGAATCCCACCCTAGAAGAAAT

SIV gag      -----
#1041         .....
              GCTGACGGCTTGTCAAGGAGTAGGGGGGCCGGGACAGAAG


SIV gag      -----
#1081         .....
              GCTAGATTAATGGCAGAAGCCCTGAAAGAGGCCCTCGCAC

SIV gag      -----
#1121         .....
              CAGTGCCAATCCCTTTTGCAGCAGCCCAACAGAGGGGACC

SIV gag      -----
#1161         .....
              AAGAAAGCCAATTAAGTGTTGGAATTGTGGGAAAGAGGGA
```


15/43

```
SIV gag -----  
#1201 .....  
      CACTCTGCAAGGCAATGCAGAGCCCCAAGAAGACAGGGAT  
  
SIV gag -----  
#1241 .....  
      GCTGGAAATGTGGAAAAATGGACCATGTTATGGCCAAATG  
  
SIV gag -----  
#1281 .....  
      CCCAGACAGACAGGCGGGTTTTTTAGGCCTTGGTCCATGG  
  
SIV gag -----  
#1321 .....  
      GGAAAGAAGCCCCGCAATTTCCCATGGCTCAAGTGCATC  
  
SIV gag -----  
#1361 .....  
      AGGGGCTGATGCCAACTGCTCCCCCAGAGGACCCAGCTGT  
  
SIV gag -----  
#1401 .....  
      GGATCTGCTAAAGAACTACATGCAGTTGGGCAAGCAGCAG  
  
SIV gag -----  
#1441 .....  
      AGAGAAAAGCAGAGAGAAAGCAGAGAGAAGCCTTACAAGG  
  
SIV gag -----  
#1481 .....  
      AGGTGACAGAGGATTTGCTGCACCTCAATTCTCTTTGG  
  
SIV gag -----  
#1521 .....  
      AGGAGACCAGTAG
```

**FIGURE 5**


17/43

FIGURE 6




18/43

FIGURE 7



19/43

FIGURE 8



20/43

FIGURE 9

1 CCTGGCCATT GCATACGTTG TATCCATATC ATAATATGTA CATTTATATT GGCTCATGTC CAACATTACC  
 72 GCCATGTTGA CATTGATTAT TGACTAGTGA TTAATAGTAA TCAATTACGG GGTCAATTAGT TCATAGCCCA  
 141 TATATGGAGT TCCGCGTTAC ATAACCTTACG GTAAATGGCC CGCCTGGCTG ACCGCCCAAC GACCCCGGCC  
 211 CATTGACGTC AATAATGACG TATGTTCCCA TAGTAACGCC AATAGGGACT TTCCATTGAC GTCAATGGGT  
 281 GGAGTATTTA CGGTAAACTG CCCACTTGGC AGTACATCAA GTGTATCATA TGCCAACTAC GCCCCCTATT  
 351 GACGTCAAATG ACGGTAAATG GCGCGCTGGC CATTTATGCCC AGTACATGAC CTTATGGGAC TTTCCCTACTT  
 421 GGCAGTACAT CTACGTATTA GTCATCGCTA TTACCATGGT GATGCGGTTT TGGCACTACA TCAATGGGCG  
 491 TGGATAGCGG TTGACTCAG GGGGATTTCG AAGTCTCCAC CCCATTGACG TCAATGGGAG TTGTGTTTGG  
 561 CACCAAAATC AACGGGACTT TCCAAAATGT CGTAACAACCT CGGCCCATTT GACGCAATG GCGCGTAGGC  
 631 GTGTACGGTG GGAGGTCTAT ATAAGCAGAG CTCGTTTAST GAACCGTCAG ATCGCCTGGA GACGCCATCC

Sall (758)

701 ACGCTGTTT GACCTCCATA GAAGACACCG GGACCGATCC AGCCTCCGCG GGCGCGCGTC GACAGAGAGA  
 771 TGGGTGCGAG AGCGTCACTA TTAAGCGGGG GAGAAATTAGA TCGATGGGAA AAAATTCGGT TAAGGCCAGG  
 841 GGGAAAGGAG AAGTACAGC TAAAGCACAT CGTATGGGCA AGCAGGGAGC TAGAAGCAAT CGCAGTTAAT  
 911 CCTGGCCTGT TAGAAACATC AGAAGGCTGT AGACAATAC TGGGACAGCT ACAACCATCC CTTACAGACG  
 981 GATCAGAGGA GCTTCGATCA CTATACAACA CAGTAGCAAC CCTCTATTGT GTGCCACGCG GGATCGAGAT  
 1051 CAAGGACACC AAGGAAGCTT TAGACAAGAT AGAGGAAGAG CAAAACAAGT CCAAGAGAGG GCGCCAGCAG  
 1121 GCAGCAGCTG ACACAGGACA CAGCAATCAG GTCAGCCAAA ATTACCTTAT AGTGACAGAC ATCCAGGGGC  
 1191 AAATGGTACA TCAGGCCATA TCACCTAGAA CTTTAAATGC ATGGGTAAAA GTAGTAGAAG AGAAGGCTTT  
 1261 CAGCCGAGAA GTGATACCCA TGTTTTCAGC ATTATCAGAA GGAGCCACCC CACAGGACCT GAACACGATG  
 1331 TTGAACACCG TGGGGGGACA TCAAGCAGCC ATGCAATGT TAAAGAGAGC CATCAATGAG GAAGCTGCAG  
 1401 AATGGGATAG AGTGCAATCA GTGCATGAG GGCCTATTGC ACCAGGCCAG ATGAGAGAAC CAAGGGGAG  
 1471 TGACATAGCA GGAACCTACTA GTACCTTCA GGAACAAATA GGATGGATGA CAAATAATCC ACCTATCCCA  
 1541 GTAGGAGAGA TCTACAAGAG GTGGATAATC CTGGGATTGA ACAAGATCGT GAGGATGTAT AGCCCTACCA  
 1611 GCATTCTGGA CATAAGACAA GGACCAAGG AACCTTTAG AGACTATGTA GACCGTTCT ATAAACTCT  
 1681 AAGAGCTGAG CAAGCTTAC AGGAGGTAA AAATTGGATG ACAGAAACCT TGTGTTCCA AAATGCGAAC  
 1751 CCAGATTGTA AGACCATCCT GAAGGCTCTC GCGCCAGCGG CTACACTAGA AGAAATGATG ACAGCATGTC  
 1821 AGGGATAGG AGGACCCGGC CATAAGGCAA GAGTTTGGC CGAGGCGATG AGCCAGGTGA CGAACTCGGC  
 1891 GACCATAATG ATGCAGAGAG GCACTTCCG GAACCGCGG AAGATCGTCA AGTGCTTCAA TTGTGGCAA



1961 GAAGGGCACA CCGCCAGGAA CTGCCGGGCC CCCCGGAAGA AGGGCTGTTG GAAATGTGGA AAGGAAGGAC  
2031 ACCAAATGAA AGATTGTACT GAGAGACAGG CTAATTTTTT AGGGAAGATC TGGCCTTCCT ACAAGGGAAG  
2101 GCCAGGGAAT TTTCTTCAGA GCAGACCAGA GCCAACAGCC CCACCAGAAG AGAGCTTCAG GTCTGGGGTA  
2171 GAGACAACAA CTCCCCCTCA GAAGCAGGAG CCGATAGACA AGGAAGTGTG TCCCTTAACT TCCCTCAGAT  
2241 CACTCTTTGG CAACGACCCC TCGTCACAGT AAGGATCGGG GGGCAACTCA AGGAAGCGCT GCTCGATACA  
2311 GGAGCAGATG ATACAGTATT AGAAGAAATG AGTTTGCCAG GAAGATGGAA ACCAAAAATG ATAGGGGGGA  
2381 TCGGGGGCTT CATCAAGGTG AGGCAGTACG ACCAGATACT CATAGAATC TGTGGACATA AAGCTATAGG  
2451 TACAGTATTA GTAGGACCTA CACCTGTCAA CATAATTGGA AGAAATCTGT TGACCCAGAT CGGCTGCACC  
2521 TTGAACCTCC CCATCAGCCC TATTGAGACG GTGCCCCGTA AGTTGAAGCC GGGGATGGAC GGCCCCAAGG  
2591 TCAAGCAATG GCCATTGACG AAGAGAAGA TCAAGGCTTT AGTCGAAATC TGTACAGAGA TGGAGAAGGA  
2661 AGGGAAGATC AGCAAGATCG GGCCTGAGAA CCCCTACAAC ACTCCAGTCT TCGCAATCAA GAAGAAGGAC  
2731 AGTACCAAGT GGAGAAAGCT GGTGGACTTC AGAGAGCTGA ACAAGAGAAC TCAGSACTTC TGGGAAGTTC  
2801 AGCTGGGCAT CCCACATCCC GCTGGGTTGA AGAAGAAGAA GTGCTGACA GTGCTGGATG TGGGTGATGC  
2871 CTACTCTCC GTTCCCTTGG ACGAGGACTT CAGGAAGTAC ACTGCCCTCA CGATACCTAG CATCAACAAC  
2941 GAGACACCAG GCATCCGCTA CCAGTACAAC GTGCTGCCAC AGGGATGGAA GGGATCACA GCCATCTTTC  
3011 AAGCAGCAT GACCAAGATC CTGGAGCCCT TCCGCAAGCA AAACCCAGAC ATCGTGATCT ATCAGTACAT  
3081 GGACGACCTC TACGTAGGAA GTGACCTGGA GATCGGSCAG CACAGGACCA AGATCGAGGA GCTGAGACAG  
3151 CATCTGTTGA GGTGGGACT GACCACACCA GACAAGAAGC ACCAGAAGGA ACCTCCCTTC CTGTGGATGG  
3221 GCTACGAAT GCATCCGAC AAGTGSACAG TGCAGCCCAT CGTGCTGCTT GAGAAGGACA GCTGGACTGT  
3291 GAACGACATA CAGAAGCTCG TGGGCAAGTT GAAGTGGSCA AGCCAGATCT ACCCAGGCAT CAAAGTTAGG  
3361 CAGCTGTGCA AGCTGCTTCG AGGAACCAAG GCACTGACAG AAGTGATCCC ACTGACAGAG GAAGCAGAGC  
3431 TAGAACTGGC AGAGAACCGA GAGATCCTGA AGGAGCCAGT ACATGGAGTG TACTACGACC CAAGCAAGGA  
3501 CCTGATCGCA GAGATCCAGA AGCAGGGGCA AGGCCAATGG ACCTACCAA TCTACCAGGA GCCCTTCAAG  
3571 AACCTGAAGA CAGGCAAGTA CGCAAGGATG AGGGGTGCCC ACACCAACGA TGTGAAGCAG CTGACAGAGG  
3641 CAGTGCAAGG GATCACCACA GAGAGCATCG TGATCTGGGG CAAGACTCCC AAGTTCAAGC TGCCCATACA  
3711 GAAGGAGACA TGGGAGACAT GGTGGACCGA GTACTGGCAA GCCACTTGA TCCCTGAGTG GGAGTTCTGT  
3781 AACACCCCTC CTTTGGTGAA ACTGTGGTAT CAGCTGGAGA AGGAACCCAT CGTGGGAGCA GAGACCTTCT  
3851 AGGTGATGG GGCAGCCAAC AGGGAGACCA AGCTGGGCAA GGCAGGCTAC GTGACCAACC GAGGACGACA  
3921 GAAAGTGGTG ACCCTGACTG ACACCACCAA CCAGAAGACT GAGCTGCAAG CCATCTACCT AGCTCTGCAA  
3991 GACAGCGGAC TGGAAGTGAA CATCTGACA GACTCACAGT ACGCACTGGG CATCATCCAA GCACAACCAG

22/43

4061 ACCAATCCGA GTCAGAGCTG GTGAACCAGA TCATCGAGCA GCTGATCAAG AAGGAGAAAG TGTACCTGGC  
4131 ATGGGTACCA GCACACAAAG GAATTGGAGG AATGAACAA GTAGATAAAT TAGTCAGTGC TGGGATCCGG  
4201 AAGGTGCTGT TCCTGGACGG GATCGATAAG GCCCAAGATG AACATGAGAA GTACCACTCC AACTGGGGGG  
4271 CTATGGCCAG CGACTTCAAC CTGCCACCTG TAGTAGCAAA AGAAATAGTA GCCAGCTGTG ATAAATGTCA  
4341 GCTAAAAGGA GAAGCCATGC ATGGACAAGT AGACTGTAGT CCAGGAATAT GGCAGCTGGA CTGCACGCAC  
4411 CTGGAGGGGA AGGTGATCCT GGTAGCAGTT CATGTAGCCA GTGGATATAT AGAAGCAGAA GTTATCCCTG  
4481 CTGAACTGG GCAGGAAACA GCATATTTC TTTTAAAT AGCAGGAAGA TGGCCAGTAA AAACAATACA  
4551 CACGGACAAC GGAAGCAACT TCACTGGTGC TACGGTTAAG GCCGCTGTT GGTGGGGGG AATCAAGCAG  
4621 GAATTGGAA TTCCCTACAA TCCCCAATCG CAGGAGTGC TGGAGAGCAT GAACAAGGAG CTGAAGAAGA  
4691 TCATCGGACA AGTGAGGGAT CAGGCTGAGC ACCTGAAGAC AGCAGTGCAG ATGGCAGTGT TCATCCACAA  
4761 CTTCAAAGA AAAGGGGGGA TTGGGGGGTA CAGTGCAGGG GAAAGGATCG TGGACATCAT GCCCACCAGC  
4831 ATCCAAACCA AGGAGCTGCA GAAGCAGATC ACCAAGATCC AGAACTTCCG GGTGTACTAC CGCGACAGCC  
4901 GCAACCCACT GTGGAAGGGA CCAGCAAAGC TCCTCTGGAA GGGAGAGGGG GCAGTGGTGA TCCAGGACAA  
4971 CAGTGACATC AAAGTGGTGC CAAGGCGCAA GGCCAAGATC ATCCGCGACT ATGGAAACA GATGGCAGGT  
5041 GATGATTGTG TGGCAAGTAG ACAGGATGAG GATTAGAACC TGGAAGAGCC TGGTGAAGCA CCATATGGCG

NheI (5117)

BstBI (5111)

5111 TTCGAAGCTA GCCTCGAGAT CCAGACTGC TGTGCTTCT AGTGGCAGC CATCTGTCT TTGCCCCCTC  
5181 CCCGTGCTT CCTTGACCT GGAAGGTGCC ACTCCCACTG TCTTTCTTA ATAAATGAG GAAATTGCAT  
5251 CGCATTGTCT GAGTAGCTGT CATCTATTC TGGGGGCTG GGTGGGCGAG CACAGCAAGG GGGAGGATTG  
5321 GGAAGACAAT AGCAGGCATG CTGGGATGC GGTGGGCTCT ATGGTACTC AGGTGCTGAA GAATTGACCC  
5391 GGTCTCTCT GGGCCAGAAA GAAGCAGGCA CATCCCTTC TCTTGACAC ACCCTGTCCA CGCCCCCTGT  
5461 TCTTAGTTC AGCCCACTC ATAGGACACT CATAGCTCAG GAGGCTCCG CCTTCAATCC CACCCGCTAA  
5531 AGTACTTGA GCGGTCTCT CCTCCCTCAT CAGCCATCA AACCAACCT AGCCTCCAG AGTGGGAAGA  
5601 AATTAAAGCA AGATAGGCTA TTAAGTGCAG AGGAGAGAA AATCCCTCCA ACATGTGAGG AAGTAATGAG  
5671 AGAATCATA GAATTTCCTC CGCTTCTCTG CTCACTGACT CGCTCGCTC GGTCTGTCGG CTGCGGGGAG  
5741 CGGTATCAGC TCACTCAAAG GCGGTAATAC GGTATCCAC AGAATCAGG GATAACGCAG GAAAGAACAT  
5811 GTGAGCAAAA GGCCAGCAAA AGGCCAGGAA CGTAAAGAG GCCCGGTTGC TGGCGTTTTT CCATAGGCTC  
5881 CGCCCCCTG ACGAGCATCA CAAAAATCGA CGCTCAAGTC AGAGGTGGCG AAACCCGACA GGACTATAAA  
5951 GATACCAGGC GTTTCCTCTT GGAAGCTCCC TCGTGGCTC TCTGTTCGG ACCCTGCGGC TTACCGGATA  
6021 CCTGTCCGCC TTCTCTCTT CCGGAAGCGT GCGGCTTCT CAATGCTCAC GCTGTAGGTA TCTCAGTTCC  
6091 GTGTAGGTCG TTGCTCCAA GCTGGGCTGT GTGCACGAAC CCCCCTTCA GCCCGACCGC TGGCCCTTAT  
6161 CCGGTAACCTA TCGTCTTGAG TCCAACCCGG TAAGACACGA CTTATCGCCA CTGGCAGCAG CCACTGGTAA  
6231 CAGGATTAGC AGAGCGAGGT ATGTAGGCGG TGCTACAGAG TTCTTGAAGT GGTGGCTAA CTACGGCTAC

23/43

6301 ACTAGAAGGA CAGTATTTGG TATCTGCGCT CTGCTGAAGC CAGTTACCTT CGGAAAAAGA GTTGGTAGCT  
6371 CTTGATCCGG CAAACAAACC ACCGCTGGTA GCGGTGGTTT TTTTGTTCG AAGCAGCAGA TTACCGGCAG  
6441 AAAAAAAGGA TCTCAAGAAG ATCCTTTGAT CTTTTCACG GGGTCTGACG CTCAGTGGAA CGAAAACTCA  
6511 CGTTAAGGGA TTTTGGTCAT GAGATTATCA AAAAGGATCT TCACCTAGAT CCTTTTAAAT TAAAAATGAA  
6581 GTTTTAAATC AATCTAAAGT ATATATGAGT AAACCTGGTC TGACAGTTAC CAATGCTTAA TCAGTGAGGC  
6651 ACCTATCTCA GCGATCTGTC TATTTGCTTC ATCCATAGTT GCCTGACTCC GGGGGGGGGG GCGCGTGAGG  
6721 TCTGCCTCGT GAAGAAGGTG TTGCTGACTC ATACCAGGCC TGAATCGCCC CATCATCCAG CCAGAAAGTG  
6791 AGGGAGCCAC GGTGTATGAG AGCTTTGTGT TAGGTGGACC AGTTGGTGAT TTTGAACCTT TGCTTTGCCA  
6861 CGGAACGGTC TGGCTTGTGG GGAAGATGCG TGATCTGATC CTTCAACTCA GCAAAAGTTC GATTTATTC  
6931 ACAAAGCCGC CGTCCCGTCA AGTCAGCGTA ATGCTCTGCC AGTGTTACAA CCAATTAACC AATTCTGATT  
7001 AGAAAACTC ATCGAGCATC AAATGAAACT GCAATTATT CATATCAGGA TTATCAATAC CATATTTTGT  
271<sup>4</sup> PhePheGlu AspLeuMetL euHisPheGlu nLeuLysAsn MetAspProA snAspIleGly TyrLysGln  
7071 AAAAAAGCCGT TTCTGTAATG AAGGAGAAA CTCACCGAGG CAGTTCCATA GATGGCAAG ATCCTGGTAT  
248<sup>4</sup> PheLeuArgL ysGlnLeuSe rProSerPhe GluGlyLeuC ysAsnTrpLe uIleAlaLeu AspGlnTyrA  
7141 CGGTCTCGGA TTCCGACTCG TCCAACATCA ATACAACCTA TTAATTTCCC CTCGTCAAAA ATAAGGTTAT  
224<sup>4</sup> rgAspAlaIle GlyValArg GlyValAspIle CysGlyIleLeuLysGly GluAspPheIleLeuAsnAs  
7211 CAAGTGAGAA ATCACCATGA GTGACGACTG AATCCGGTGA GAATGGCAAA AGCTTATGCA TTTCTTTCCA  
201<sup>4</sup> pLeuSerPhe AspGlyHisT hrValValSe rAspProSer PheProLeuL euLysHisMe lGluLysTrp  
7281 GACTTGTTC A CAGGCCAGC CATTACGCTC GTCATCAAAA TCACCTCGCAT CAACCAAAACC GTTATTTCATT  
178<sup>4</sup> ValGlnGluV alProTrpGly AsnArgGlu AspAspPheA spSerAlaAs pValLeuGly AsnAsnMetA  
7351 CGTGATTGCG CCTGAGCGAG ACGAAATACG CGATCGCTGT TAAAAGGACA ATTACAAACA GGAATCGAAT  
154<sup>4</sup> rgSerGlnAlaGlnAlaLeu ArgPheValA rgAspSerAs nPheProCys AsnCysValP rolleSerHi  
7421 GCAACCGCG CAGGAACACT GCCAGCGCAT CAACAATATT TTCACCTGAA TCAGGATATT CTCTAATAC  
131<sup>4</sup> sLeuArgArg LeuPheValA laLeuAlaAs pValIleAsn GluGlySerA spProTyrGly uGluLeuVal  
7491 CTGGAATGCT GTTTTCCCGG GGATCGCAGT GGTGAGTAAC CATGCATCAT CAGGAGTACG GATAAAATGC  
108<sup>4</sup> GlnPheAlaT hrLysGlyPr olleAlaThr ThrLeuLeuT rpAlaAspAs pProThrArg llePheHisL  
7561 TTGATGGTGG GAAGAGGCAT AAATTCCGTC AGCCAGTTTA GTCTGACCAT CTCATCTGTA ACATCATGG  
84<sup>4</sup> ysIleThrPr oLeuProMet PheGluThrL euTrpAsnLe uArgValMet GluAspThrV alAspAsnAl  
7631 CAACGCTACC TTTGCCATGT TTCAGAAACA ACTCTGGCGC ATCGGGCTTC CCATACAATC GATAGATTGT  
61<sup>4</sup> aValSerGly LysGlyHisL ysLeuPheLe uGluProAla AspProLysG lyTyrLeuAr gTyrIleThr  
7701 CGCACCTGAT TGCCCGACAT TATCGCGAGC CCATTTATAC CCATATAAAT CAGCATCCAT GTTGGAAATTT  
38<sup>4</sup> AlaGlySerG lnGlyValAs nAspArgAla TrpLysTyrG lyTyrLeuAs pAlaAspMet AsnSerAsnL  
7771 AATCGCGGCC TCGAGCAAGA CGTTTCCCGT TGAATATGGC TCATAACACC CCTTGTATTA CTGTTTATGT  
14<sup>4</sup> euArgProAr gSerCysSer ThrGluArgG lnIleHisSe rMet  
7841 AAGCAGACAG TTTTATGTGT CATGATGATA TATTTTATC TTGTGCAATG TAACATCAGA GATTTTGAGA  
7911 CACAACGTGG CTTTCCCCC CCCCCATTA TTGAAGCATT TATCAGGGTT ATTGTCTCAT GAGCGGATAC  
7981 ATATTTGAAT GTATTTAGAA AAATAAACAA ATAGGGGTTT CGCGCACATT TCCCCGAAAA GTGCCACCTG  
8051 ACGTCTAAGA AACCATTTATT ATCATGACAT TAACCTATAA AAATAGGCGT ATCACGAGGC CCTTTCTGCT  
8121 CGCGCGTTTC GGTGATGACG GTGAAAACCT CTGACACATG CAGCTCCCGG AGACGGTCCAG AGCTTGCTCTG  
8191 TAAGCGGATG CCGGGAGCAG ACAAGCCCGT CAGGGCGCGT CAGCGGGTGT TGGCGGGTGT CCGGGCTGGC  
8261 TTAACATATGC GGCATCAGAG CAGATTGTAC TGAGAGTGCA CCATATGCGG TGTGAAATAC CGCACAGATG  
8331 CGTAAGGAGA AAATACCGCA TCAGATTGGC TATTGG

24/43

FIGURE 10

1 TGGAAGGGCT AATTGTGTC CAAAAAGAC AAGAGATCCT TGATCTGTGG ATCTACCACA CACAAGGCTA  
 71 CTTCCTTGAT TGGCAGAACT ACACACCAGG GCCAGGGATC AGATATCCAC TGACCTTTGG ATGGTGCTTC  
 141 AAGTTAGTAC CAGTTGAACC AGAGCAAGTA GAAGAGGCCA AATAAGGAGA GAAGAACAGC TTGTTACACC  
 211 CTATGAGCCA GCATGGGATG GAGGACCCGG AGGAGAGAAT ATTAGTGTGG AAGTTTGACA GCTTCCTAGC  
 281 ATTTCTGCAC ATGGCCCCGAG AGCTGCATCC GGAGTACTAC AAAGACTGCT GACATCGAGC TTTCTACAAG  
 351 GGACTTTCCG CTGGGGACTT TCCAGGGAGS TGTGGCCTGG GCGGGACTGG GAGTGGCGA GCTTCAGAT  
 421 GCTACATATA AGCAGCTGCT TTTTGCTGT ACTGGGCTTC TCTGGTTAGA CCAGATCTGA GCTTGGGAGC  
 491 TCTCTGGCTA ACTAGGGAAC CCACTGCTTA AGCCTCAATA AAGCTTGCCT TGAGTGCTCA AAGTAGTGTG  
 561 TGCCCGTCTG TTGTGTGACT CTGGTAACTA GAGATCCCTC AGACCCCTTT AGTCAGTGTG GAAATCTCT  
 631 AGCAGTGGCG CCCGAACAGG GACTTGAAG CGAAAGTAAA GCCAGAGGAG ATCTCTCGAC GCAGGACTCG  
 BssHII (711)  
 701 GCTTGCTGAA GCGGCGacgg caapaggoga ggggggggc cggacggga ggggaaaaa ttgactagg  
 Clal (830)  
 771 gggggctaga aggagagagc tgggtgggag agcctcagta ttgaggggg gaggattaga tggatgggaa  
 841 AAAATTCGGT TAAGGCCAGG GGGAAAGAAA AAATATAAT TAAATATAT AGTATGGCA AGCAGGGAGC  
 AccI (959)  
 911 TAAAGCGATT CGCAGTTAAT COTGGCTGT TAGAAGATC AGAAGGCTGT AGACAAATAC TGGGACAGCT  
 981 ACAACCATCC CTTGAGACAG GATGAGAGA ACTTAGATCA TTATATATA CAGTAGCAAC CCTCTATTGT  
 1051 GTGCATCAA GGTAGAGAT AAAAGACACC AAGGAGCTT TAGACAAGAT AGAGGAAGAG CAAAACAAA  
 1121 GTAAGAAAA AGCACAGCAA GCAGCAGCTG ACACAGGACA CAGCAATCAG GTCAGCCAAA ATTACCCAT  
 1191 AGTGCAAAAC ATCCAGGGGC AAATGGTACA TCAGGCCATA TCACCTAGAA CTTTAAACGA TAAGCTTGGG  
 1261 AGTTCCGGGT TACATAACTT ACGSTAATG GCGCGCTGG CTGACCGCCC AACGACCCC GCCCATGAC  
 1331 GTCATAATG ACGTATGTC CCAATGTAAC GCCAATAGG ACTTTCATT GACGTCAATG GGTGGAGTAT  
 1401 TTACGGTAAA CTGCCCACTT GGCAGTACAT CAAGTGTATC ATATGCCAAG TACGCCCCCT ATTGACGTCA  
 1471 ATGACGGTAA ATGGCCCGCC TGGCATTATG CCCAGTACAT GACCTTATGG GACTTTCCTA CTGGGCAGTA  
 1541 CATCTACGTA TTAGTCATCG CTATTACCAT GGTGATGCGG TTTTGGCAGT ACATCAATGG GCGTGATAG  
 1611 CGGTTTGAAT CACGGGGATT TCCAAGTCTC CACCCCATG ACGTCAATGG GAGTTTGT TTGGCACAAA  
 1681 ATCAACGGGA CTTTCCAAA TGTCGTAACA ACTCCGCCCC ATTGACGCAA ATGGGCGGTA GCGGTGTACG  
 1751 GTGGGAGGTC TATATAAGCA GAGCTCGTTT AGTGAACCGT CAGATCGCCT GGAGACGCCA TCCACGCTGT

25/43

1821 TTTGACCTCC ATAGAAGACA CCGACTCTAG ASgattccatc TAAGTAAGCT TGGCATTCCG GTACTGTTCG  
 1891 TAAATGGAA GAGGCCAAA ACATAAGAA AGGCCCGCG CCATTCTATC CTCTAGAGGA TGGAAACGGT  
 1961 GGAGAGCAAC TGCATAAGGC TATGAAGAGA TACGCCCTGG TTCTTGGAAc ANTTCCTTTT ACAGATGCAC  
 2031 ATATCGAGGT GAACATCAGG TACGCCGAAT ACTTCGAAT GTCCGTTCCG TTGGCAGAAG CTATGAAACG  
 2101 ATATGGGCTG AATACAAATC ACAGAATCGT CGTATGCAGT GAAAACTCTC TTCAATTCTT TATGCCGCTG  
 2171 TTGGGCGCGT TATTTATCGG AGTTGCAGTT GCGCCCGCGA ACGACATTTA TAATGAACGT GAATTGCTCA  
 2241 ACASTATGAA CATTTCGCAG CCTACCGTAG TGTGTGTTTC CAAAAGGGG TTGCAAAAAA TTTTGAACGT  
 2311 GCAAAAAAA TTACCAATAA TCCAGAAAT TATTATCATG GATTCTAAAA CCGATTACCA GCGATTTCAG  
 2381 TCGATGTACA CGTTCGTAC ATCTCATCTA CCTCCCGGTT TTAATGAATA CGATTTTCTA CCGAGTCTCT  
 2451 TTGATCGTGA CAAACAATT GCACTGATAA TGAATTCCTC TGGATCTACT GGGTTACCTA AGGGTGTGGC  
 2521 CCTCCCGCAT AGAATGCGCT GCGTCAGATT CTCGCATGCC AGAGATCCTA TTTTGGCAA TCAATCAATT  
 2591 CCGGATACGT CGATTTTAAG TGTGTTCCTA TTCCATCAGG GTTTTGAAT GTTACTACA CTCGGATATT  
 2661 TGATATGTGG ATTTCCAGTC GTCTTAATGT ATAGATTGA AGAAGAGCTG TTTTACGAT CCCTTCAGGA  
 2731 TTACAAAATT CAAATGCGT TGCTASTACC AACCTATTT TCATTCTTCG CCAAAAGCAC TCTGATTGAC  
 2801 AATACGATT TATCTAATTT ACACGAAAT GCTTCTGGG GCGACCTCT TTGAAAGAA GTCCGGGAAG  
 2871 CGGTGCGAAA ACGTTTCAT CTTCAGGGA TACGACAGG ATATGGGCTC ACTGAGACTA CATCAGCTAT  
 2941 TCTGATTACA CCCGAGGGG ATGATAAACC GGGGCGGTC GTTAAAGTTG TTCCATTTT TGAAGCGAAG  
 3011 GTTGTGGATC TGGATACCGG GAAAACGCTG GCGCTTAATC AGAGAGCGGA ATTATGTGTC AGAGGACCTA  
 3081 TGATTATGTC CGGTATGTA AACATCCCG AAGCGACCA CCGCTTGATT GACAAGGATG GATGGCTACA  
 3151 TTCTGGAGAC ATAGCTTACT GGGACGAGA CGACACTTC TTCTAGTTG ACCGCTTGAA GTCTTTAATT

### Clal (3259)

3221 AATACAAAG GATATCAGGT GGGCCCGCT GAATTGGAAT CGATATTGTT ACAACACCCC AACATCTTCG  
 3291 ACGCGGGCGT GCGAGGTCTT CCGAGCATG ACGCCGCTGA ACTTCCCGCC GCGGTGTGTT TTTTGGAGCA  
 3361 CGGAAAGACG ATGACGGAAA AAGAGATCGT GGAATTACGTC GCCAGTCAAG TAACACCCG GAAAAAGTTG  
 3431 CGCGGAGGAG TTGTGTTTGT GGACGAATA CCGAAGGTC TTACCGGAAA ACTCGAGCCA AGAAAAATCA

### Apal (3557)

### XhoI (3548) KpnI (3563)

3501 GAGAGATCTT CATAAAGGCC AAGAAAGGCG GAAATCCAA ATTGTAACTC GAGGGGGGGC CCGGTACCTT  
 3571 TAAGACCAAT GACTTACAAG GCAGCTGTAG ATCTTAGCCA CTTTTAAAA GAAAAGGGGG GACTGGAAGG  
 3641 GCTAATTCAC TCCCAAGAA GACAAGATAT CCTTGATCTG TGGATCTACC ACACACAAGG CTACTTCCCT

26/43

3711 GATTGGCAGA ACTACACACC AGGGCCAGGG GTCAGATATC CACTGACCTT TGGATGGTGC TACAAGCTAG  
3781 TACCAGTTGA GCCAGATAAG GTAGAAGAGG CCAATAAAGG AGAGAACACC AGCTTGTTAC ACCCTGTGAG  
3851 CCTGCATGGA ATGGATGACC CTGAGAGAGA AGTGTTAGAG TGGAGGTTTG ACAGCCGCCCT AGCATTTTCAT  
3921 CACGTGGCCC GAGAGCTGCA TCCGGAGTAC TTCAAGAACT GCTGACATCG AGCTTGCTAC AAGGGACTTT  
3991 CCGCTGGGGA CTTTCCAGGG AGGCTGGCC TGGCCGGGAC TGGGGAGTGG CGAGCCCTCA GATGCTGCAT  
4061 ATAAGCAGCT GCTTTTGGCC TGTACTGGGT CTCTCTGGTT AGACCAGATC TGAGCCTGGG AGCTCTCTGG  
4131 CTAAGTAGGG AACCCACTGC TTAAGCCCTCA ATAAAGCTTG CCTTGAGTGC TTCAAGTAGT GTGTGCCCGT  
4201 CTGTTGTGTG ACTCTGGTAA CTAGAGATCC CTCAGACCCCT TTTAGTCAGT GTGGAAAAATC TCTAGCAGCC  
4271 CCCAGGAGGT AGAGGTTGCA GTGAGCCAAG ATCGCGCCAC TGCATTCCAG CCTGGGCAAG AAAACAAGAC  
4341 TGTCTAAAAAT AATAATAATA AGTTAAGGGT ATTAAATATA TTTATACATG GAGGTCATAA AAATATATAT  
4411 ATTTTGGGCTG GGCGCAGTGG CTCACACCTG CGCCCGGCCCT TTTGGGAGGC CGAGGCAGGT GGATCACCTG  
4481 AGTTTGGGAG TTCCAGACCA GCCTGACCAA CATGGAGAAA CCCCTTCTCT GTGTATTTT AGTAGATTTT  
4551 ATTTTATGTG TATTTTATTC ACAGGTATTT CTGGAAGAACT GAAACTGTTT TTCTCTACT CTGATACCAC  
4621 AAGAATCATC AGCAGAGAGG AAGACTTCTG TGATCAAAATG TGGTGGGAGA GGGAGGTTT CACCAGCACA  
4691 TGAGCAGTCA GTTCTGCCGC AGACTCGGC GGTGTCCTTC GGTTCAGTTC CAACACCCGC TGCCTGGAGA  
4761 GAGGTCAGAC CACAGGGTGA GGGCTCAGTC CCCAAGACAT AAACACCCAA GACATAAACA CCCAACAGGT  
4831 CCACCCCGCC TGCTGCCAG GCAGAGCCGA TTCACCAAGA CGGAATTAG GATAGAGAAA GAGTAAGTCA  
4901 CACAGAGCCG GCTGTGCCGG AGAACGGAGT TCTATTATGA CTCAAATCAG TCTCCCAAG CATTCGGGGA  
4971 TCAGAGTTT TAAGGATAAC TTAGTGTGTA GGGGGCCAGT GAGTTGGAGA TGAAAGCGTA GGGAGTCGAA  
5041 GGTGTCTTT TGCGCCGAGT CAGTTCTCG GTGGGGCCA CAAGATCGGA TGAGCCAGTT TATCAATCCG  
5111 GGGTGGCCAG CTGATCCATG GAGTGCAGGG TCTGCAAAAT ATCTCAAGCA CTGATTGATC TTAGGTTTTA  
5181 CAATAGTAT GTTACCCAG GAACAATTG GGGAGGTCA GAATCTGTA GCCTGTAGCT GCATGACTCC  
5251 TAAACCATAA TTCTTTTTT GTTTTTTTTT TTTTATTTT GAGACAGGT CTCACCTGT CACCTAGGCT  
5321 GGAGTGCAGT GTGCAATCA CAGCTCAGT CAGCCCTAG AGCGCCGCC ACCCGGCTG AGCTCCAATT  
5391 CGCCCTATAG TGAGTCGTAT TACAATTCAC TGGCCGTCTG TTTACAACGT CGTGACTGGG AAAACCCCTG  
5461 CGTTACCCAA CTTAATCGCC TTGCAGCACA TCCCCCTTC GCCAGCTGGC GTAATAGCGA AGAGGCCCGC  
5531 ACCGATCGCC CTTCCCAACA GTTGGCCAGC CTGAATGGCG AATGGCGCGA AATTGTAAAC GTTAATATT  
5601 TGTTAAATTT CGCGTTAAAT TTTTGTATAA TCAGTCAAT TTTTAAACCA TAGGCCGAAA TCGGCAAAAT  
5671 CCCTTATAAA TCAAAAGAA AGACCGAGAT AGGTTGAGT GTTGTTCAG TTTGGAACAA GAGTCCACTA  
5741 TTAAGAACG TGAGTCCAA CGTCAAGAGG CGAAAAACCG TCTATCAGGG CGATGGCCCA CTACGTGAAC  
5811 CATCACCTTA ATCAAGTTTT TTGGGGTCGA GGTGCCGTA AGCACTAAAT CGGAACCTTA AAGGGAGCCC  
5881 CCGATTTAGA GCTTGACGGG GAAAGCCGGC GAAGTGCGC AGAAAGGAAG GGAAGAAAGC GAAAGGAGCG  
5951 GCGCTAGGG CGCTGGCAAG TGTAGCGGC ACGCTGCGC TAACCACCAC ACCCGCCGCG CTTAATCGCG  
6021 CGCTACAGGG CGCGTCCAG GTGGCACTTT TCGGGGAAAT GTGCGCGGAA CCCCTATTG TTTATTTTT  
6091 TAAATACATT CAATATGTA TCCGCTCATG AGACAATAAC CCTGATAAAT GCTTCAATA TATTGAAAAA  
6161 GGAAGAGTAT GAGTATTCAA CATTTCCGTG TCGCCCTTAT TCCCTTTTTT CGGCAATTT GCCTTCTCTG  
6231 TTTTGCTCAC CCAGAAACCG TGGTGAAGT AAAAGATGCT GAAAGATCAGT TGGGTGCACG AGTGGGTTAC  
6301 ATCGAATCG ATCTCAACAG CGTAAAGATC CTTGAGAGTT TTCGCCCGA AGAAGCTTT CCAATGATGA  
6371 GCACTTTTAA AGTTCTGCTA TGTGGCGCG TATTATCCCG TATTGACGCC GGGCAAGAGC AACTCGGTG  
6441 CCGCATACAC TATTCTCAGA ATGACTTGGT TGAGTACTCA CCAGTCAAG AAAAGCATCT TACGGATGGC  
6511 ATGACAGTAA GAGAATTATG CAGTGCTGCC ATAACCATGA GTGATAACAC TGCGGCCAAC TTACTTCTGA  
6581 CAACGATCGG AGGACCGAAG GAGCTAACCG CTTTTTGTCA CAACATGGGG GATCATGTAA CTCGCCCTGA  
6651 TCGTTGGGAA CCGGAGCTGA ATGAAGCCAT ACCAAACGAC GAGCGTGACA CCACGATGCC TGTAGCAATG  
6721 GCAACACGT TCGGCAAACT ATTAACCTGC GAACACTTTA CTCTAGCTTC CCGGCAACA TTAATAGACT  
6791 GGAATGAGGC GGATAAAGTT GCAAGGACC TTCTGCGCTC GGCCCTTCCG GCTGGCTGGT TTATTGCTGA  
6861 TAAATCTGGA GCGGTGAGC GTGGTCTCG CGGTATCATT GCAGCACTGG GGCAGATGG TAAGCCCTCC  
6931 CGTATCGTAG TTATCTACAC GACGGGAGT CAGGCAACTA TGGATGAACG AAATAGACAG ATCGCTGAGA  
7001 TAGGTGCTC ACTGATTAAG CATTTGTAA TGTGAGCCA AGTTTACTCA TATATACTTT AGATTGATTT  
7071 AAAACTTCAT TTTTAAATTA AAAGGATCTA GGTGAAGATC CTTTTGTATA ATCTCATGAC CAAAATCCCT  
7141 TAACTGTAGT TTTCTGTCCA CTGAGCGTCA GACCCGCTAG AAAAGATCAA AGGATCTTCT TGAGATCCTT  
7211 TTTTCTCGC CGTAATCTGC TGCTTGCAA CAAAAAAC ACCGCTACCA GCGGTGGTTT GTTTGCGGGA  
7281 TCAAGAGCTA CCAACTCTT TTCCGAAGGT AACTGGCTTC AGCAGAGCGC AGATACCAAA TACTGTCTTT

27/43

```
7351 CTAGTGTAGC CGTAGTTAGG CCACCACTTC AAGAACTCTG TAGCAACGGC TACATACCTC GCTCTGCTAA
7421 TCCGTGTACC AGTGGCTGCT GCCAGTGGCG ATAAGTCGTG TCTTACCGGG TTGGACTCAA GACGATAGTT
7491 ACCGGATAAG GCGCAGCGGT CGGGCTGAAC GGGGGGTTCC TCCACACAGC CCAGCTTGGG GCGAACGACC
7561 TACACCGAAC TGAGATACCT ACAGCGTGAG CTATGAGAAA GCGCCACGCT TCCCGAAGGG AGAAAGGCGG
7631 ACAGGTATCC GGTAAAGCGG AGGGTCGGAA CAGGAGAGCG CACGAGGGAG CTTCAGGGG GAAACGCCCTG
7701 GTATCTTTAT AGTCCCTGTC GGTTCGCCA CCTCTGACTT GAGCGTCGAT TTTTGTGATG CTCGTCAGGG
7771 GGGCGGAGCC TATGGAAAAA CGCCAGCAAC GCGGCCCTTT TACGGTTCCT GGGCTTTTC TGGCCTTTTG
7841 CTCACATGTT CTTTCCTGCG TTATCCCTCG ATTCCTGTGA TAACCGTATT ACCGCTTTTC AGTGAGCTGA
7911 TACCGCTCGC CGCAGCCGAA CGACCGAGCG CAGCGAGTCA GTGAGCGAGG AAGCGGAAGA GCGCCCAATA
7981 CGCAAACCGC CTCCTCCCGC GCGTTGGCCG ATTCAATTAAT GCAGCTGGCA CGACAGGTTT CCGACTGGA
8051 AAGCGGGCAG TGAGCGCAAC GCAATTAATG TGAGTTAGCT CACTCATTAG GCACCCACGG CTTTACACTT
8121 TATGCTTCCG GCTCGTATGT TGTGTGGAAT TGTGAGCGGA TAACAAATTTC ACACAGGAAA CAGCTATGAC
8191 CATGATTACG CCAAGCTCGG AATTAACCTT CACTAAAGGG AACAAAAGCT GCTGCAGGGT CCCTAACTGC
8261 CAAGCCCCAC AGTGTGCCCT GAGGCTGCCC CTTCCTTCTA GCGGCTGCCC CCAGTCGGCT TTGCTTTCCC
8331 TAGTTTCAGT TACTTGCGTT CAGCCAAGGT CTGAAACTAG GTGCGCACAG AGCGGTAAAG CTGCGAGAGA
8401 AAGAGACCAG CTTTACAGGG GGTTTATCAC AGTGCAACCT GACAGTCGTC AGCCTCACAG GGGGTTTATC
8471 ACATTGCACC CTGACAGTCG TCAGCCTCAC AGGGGGTTTA TCACAGTGCA CCCTTACAAAT CATTCATTT
8541 GATTCACAAT TTTTTTAGTC TCTACTGTGC CTAACCTTGT AGTTAAATTT GATCAGAGGT GTGTTCCAG
8611 AGGGGAAAAC AGTATATACA GGGTTCAGTA CTATCGCATT TCAGGCCCTC ACCTGGGTCT TGGAAATGTT
8681 CCCCCGAGGG GTGATGACTA CTTCAAGTTGG ATCTCCACAG GTCACAGTGA CACAAGATAA CCAAGACACC
8751 TCCCAAGGCT ACCACAATGG GCGGCCCTCC ACGTGACAT GGC CGGAGGA ACTGCCATGT CGGAGGTGCA
8821 AGCACACCTG CGCATCAGAG TCCTTGGTGT GGAGGGAGGG ACCAGCGCAG CTTCCAGCCA TCCACCTGAT
8891 GAACAGAAC TAGGGAAAGC CCCAGTTCTA CTTACACCAG GAAAGGC
```

28/43

FIGURE 11

1 TGGAAGGGCT AATTGGTCC CAAAAAGAC AAGAGATCCT TGATCTGTGG ATCTACCACA CACAAGGCTA  
 71 CTTCCTTGAT TGGCAGAACT ACACACCAGG GCCAGGGATC AGATATCCAC TGACCTTTGG ATGGTGCTTC  
 141 AAGTTAGTAC CAGTTGAACC AGAGCAAGTA GAAGAGGCCA AATAAGGAGA GAAGAACAGC TTGTTACACC  
 211 CTATGAGCCA GCATGGGATG GAGGACCCGG AGGGAGAAGT ATTAGTGTGG AAGTTTGACA GCCTCCTAGC  
 281 ATTTGCTCAC ATGGCCCGAG AGCTGCATCC GGAGTACTAC AAAGACTGCT GACATCGAGC TTTCTACAAG  
 351 GGACTTTCCG CTGGGGACTT TCCAGGGAGG TGTGGCCTGG GCGGGACTGG GGAATGGCGA GCCCTCAGAT  
 421 GGTACATATA AGCAGCTGCT TTTTGCCGTG ACTGGGTCTC TCTGGTTAGA CCAGATCTGA GCCTGGGAGC  
 491 TCTCTGGCTA ACTAGGGAAC CCACTGCTTA AGCCTCAATA AAGCTTGCCCT TGAGTGCTCA AAGTAGTGTG  
 561 TGCCCGTCTG TTGTGTGACT CTGTAACATA GAGATCCCTC AGACCCCTTT AGTCAGTGTG GAAAATCTCT  
 631 AGCAGTGGCG CCCGAACAGG GACTTGAAG CGAAAGTAA GCCAGAGGAG ATCTCTCGAC GCAGGACTCG  
 BssHII (711)  
 701 GCTTGCTGAA GGGGACCTT CAGGAGGCGA GGGGAGGCG CAGAGGAGG CAGGAGGAG CAGGAGGAG  
 Clal (830)  
 771 GAGGAGGAG GAGGAGGAG TGGTGGGAG AGGTCAGTA TTAGGAGGAG GAGGAGGAG TCGATGGGAA  
 841 AAAATTCGCT TAGGAGGAG GGGAGGAG AGTACAGC TTAGGAGAT COTATGGGA ACCAGGGAGC  
 AccI (959)  
 911 TAGACGATT CCGAGTTAAT CTGGGCTGT TAGAACATC AGAGGCTGT AGACAAATAC TGGGACAGCT  
 981 ACAACCATCC CTTGAGACAG GATCAGAGG GATTGATCA CTATACACA CAGTAGCAAC CCTCTATTGT  
 1051 GTGACCCAGC GGATCGAGAT CAGGACACC AAGGAAGCTT TAGACAAGAT AGAGGAAGAG CAAAACAAGT  
 1121 CCAAGAGGAA GGGCCAGCAG GCAGCAGTG ACACAGGACA CAGCAATCAG GTCAGGCCAA ATTACCTAT  
 1191 AATGAGAAC ATCCAGGGGC AAATGGTACA TCAGGCCATA TCACCTAGAA CTTTAAACGA TAAGCTTGGG  
 1261 AGTTCGCGT TACATAACTT ACGTAAATG GCGCGCTGG CTGACCGCCC AAGGACCCCC GCCCATTTGAC  
 1331 GTCATAATG ACGTATGTC CCATAGTAAC GCCATAGGG ACTTTCATT GACGTCAATG GGTGGAGTAT  
 1401 TTACGGTAAA CTGCCACTT GGCAGTACAT CAAGTGTATC ATATGCCAAG TACGCCCCCT ATTGACGTCA  
 1471 ATGACGGTAA ATGGCCCGCC TGGCATTATG CCCAGTACAT GACCTTATGG GACTTTCCTA CTGGGAGTA  
 1541 CATCTACGTA TTAGTCATCG CTATTACCAT GGTGATCGG TTTTGGCAGT ACATCAATGS GGTGGGATAG



29/43

1611 CGGTTTGACT CACGGGGATT TCCAAGTCTC CACCCCATTC AGGTCAATGG GAGTTTGTCT TGGCACCAAA  
 1681 ATCAACGGGA CTTCACAAA TGTCGTAAAC ACTCGGCCCC ATTGACGCAA ATGGAGCGTA GCGGTGTACG  
 1751 GTGGGAGGTC TATATAAGCA GAGCTCGTTT AGTGAACCGT CAGATCGGCT GGAGACGCCA TCCACGCTGT  
 1821 TTGACCTCC ATAGAAGACA CCGACTCTAG AGGATCCATC TAAGTAAAGT TGGCATTCGG GTACTGTTCG  
 1891 TAAAATGGAA GACGCCAAAA ACATAAGAA AGGCCCGGCG CCATTCCTATC CTCTAGAGGA TGGAAACGCT  
 1961 GGAGAGCAAC TGCATAAGGC TATGAAGAGA TACGCCCTGG TTCTCGAAC AATTGCTTTT ACAGATGCAC  
 2031 ATATCGAGGT GAACATCAGC TACGCGGAAT ACTTCGAAAT GTCCGTTCGG TTGGCAGAG CTATGAACG  
 2101 ATATGGGCTG AATACAAATC ACAGAATCGT CGTATGCAAT GAAACTCTC TTCAATCTCT TATGCCGGTG  
 2171 TTGGGCGCGT TATTTATCGG AGTTGCAATT GCGCCCGCGA ACGACATTTA TAATGAACGT GAATTGCTCA  
 2241 ACAGTATGAA CATTTCGCAG CCTACCGTAG TGTTGTCTC CAAAAGGCGG TTGCAAAAA TTTTGAACGT  
 2311 GCAAAAAA TTACCAATAA TCCAGAAAAT TATTATCATG GATTCTAAAA CGGATTACCA GGGATTTACG  
 2381 TCGATGTACA CGTTTCGCAC ATCTCATCTA CCTCCCGGTT TTAATGAATA CGATTCTGTA CCAGAGTCTT  
 2451 TTGATCGTGA CAAACAATT GCACGTATAA TGAATTCCTC TGGATCTACT GGGTTACCTA AGGGTGTGGC  
 2521 CCTCCCGCAT AGAAGTGCCT GCGTCAGATT CTGCGATGCC AGAGATCCTA TTTTGGGCAA TCAATCATT  
 2591 CCGGATACCT CGATTCTAAG TGTGTGCA TTCCATCAGG GTTTTGGAA GTTTACTACA CTCGGATATT  
 2661 TGATATGTTG APTTCGATC GTCTTAATGT ATAGATTGTA AGAGAGGCTG TTTTACGAT CCCTTCAGGA  
 2731 TTACAAAAAT CAAAGTGCCT TGCTASTACC AACCTATTT TCATCTCTCG CAAAAGCAC TCTGATTGAC  
 2801 AATACGATT TATCTAATTT ACACGAATT GCTCTGCGG GCGCACCTCT TTGGAAGAA GTCCGGGAAG  
 2871 CGGTTGCAA ACGCTTCAT CTTCAGGGA TACGACAGG ATATGGGCTC ACTGAGACTA CATCAGCTAT  
 2941 TCTGATTACA CCGAGGGGG ATGATAAAC GGGCGCGGTC GGTAAAGTTG TTCCATTTT TGAAGCGAAG  
 3011 GTGTGGATC TGATACCGG GAATACGCTG GCGCTTAATC AGAGAGGCGA ATTATGTGTC AGAGGACCTA  
 3081 TGATTATGTC CGTTATGTA AACATCCGG AAGCGACCA CGCCTTGATT GACAAGGATG GATGGCTACA  
 3151 TTCTGGAGAC ATAGCTTACT GGGACGAGA CGAACCTTC TTCTAGTTG ACCGTTTGA GTCTTTAATT

# Clal (3259)

3221 AAATCAAG GATATCAGGT GCGCCCGCT GAATTGGAAT CGATATTGTT ACAACACCCC AACATCTTCG  
 3291 AGCGGGGCGT GCGAGGTCTT CCGACGATG ACGCGGTA ACTTCCCGCC GCGGTGTTC TTTGGAGCA  
 3361 CGAAAGACG ATGACGAAA AAGAGATCGT GGAATACGTC GCCAGTCAAG TAACACCGC GAAAAGTTG  
 3431 CCGGAGGAG TTGTGTTGT GGACGAAGTA CCGAAGGTC TTACCGAAA ACTCGACCA AGAAAAATCA

30/43

ApaI (3557)

XhoI (3548) KpnI (3563)

```

3501 GAGAGATCCT CATTAAGGCC AAGAAGGGCG GAAAGTCCAA ATTGTAACTC GAGGGGGGGC CCGGTACCTT
3571 TAAGACCAAT GACTTACAAG GCAGCTGTAG ATCTTAGCCA CTTTTTAAAA GAAAAGGGGG GACTGGAAGG
3641 GCTAATTCAC TCCCAAAGAA GACAAGATAT CCTTGATCTG TGGATCTACC ACACACAAGG CTACTTCCCT
3711 GATTGGCAGA ACTACACACC AGGGCCAGGG GTCAGATAAT CACTGACCTT TGGATGGTGC TACAAGCTAG
3781 TACCAGTTGA GCCAGATAAG GTAGAAGAGG CCAATAAAGG AGAGAACACC AGCTTGTATC ACCCTGTGAG
3851 CCTGCATGGA ATGGATGACC CTGAGAGAGA AGTGTTAGAG TGGAGGTTTG ACAGCCGCTT AGCATTTCAT
3921 CACGTGGCCC GAGAGCTGCA TCCGGAGTAC TTCAAGAACT GCTGACATCG ACCTTGCTAC AAGGGACTTT
3991 CCGCTGGGGA CTTTCCAGGG AGGCGTGGCC TGGGCGGGAC TGGGGAGTGG CGAGCCCTCA GATGCTGCAT
4061 ATAAGCAGCT GCTTTTGGCC TGTACTGGGT CTCTCTGGTT AGACCAGATC TGAGCCTGGG AGCTCTCTGG
4131 CTAAGTAGGG AACCCACTGC TTAAGCCTCA ATAAAGCTTG CCTTGAGTGC TTCAAGTAGT GTGTGCCCGT
4201 CTGTTGTGTG ACTCTGTGTA CTAGAGATCC CTCAGACCTT TTTAGTCAGT GTGGAAAATC TCTAGCAGCC
4271 CCCAGGAGGT AGAGGTTCGA GTGAGCCAAG ATCGCGCCAC TGCATTCCAG CCTGGGCAAG AAAACAAGAC
4341 TGTCATAAAT AATAATAATA AGTTAAGGGT ATTAATAATA TTTATACATG GAGGTCATAA AAATATATAT
4411 ATTTGGGCTG GCGCGAGTGG CTCACACCTG CGCCCGGCCC TTTGGGAGGC CGAGGCAGGT GGATCACCTG
4481 AGTTTGGGAG TTCCAGACCA GCTGACCAA CATGGAGAAA CCCCTTCTCT GTGTATTTT AGTAGATTTT
4551 ATTTTATGTG TATTTTATTC ACAGGTATTT CTGGAAGAACT GAAACTGTTT TTCTCTACT CTGATACCAC
4621 AAGAATCATC AGCAGAGAGG AAGACTTCTG TGATCAATG TGGTGGGAGA GGGAGGTTTT CACCAGCACA
4691 TGAGCAGTCA GTTCTGCCGC AGACTCGGGG GGTGTCTTTC GGTTCAGTTC CAACACCGCC TGCCCTGGAGA
4761 GAGGTGAGAC CACAGGGTGA GGGCTCAGTC CCAAGACAT AAACACCCAA GACATAAACA CCAACACAGT
4831 CCACCCGCCC TGCTGCCGAG GCAGAGCCGA TTCACCAAGA CGGGAATTAG GATAGAGAAA GAGTAAGTCA
4901 CACAGAGCCG GCTGTGCCGG AGAACGGAGT TCTATTATGA CTCAAATCAG TCTCCCAAG CATTCGGGGA
4971 TCAGAGTTTT TAAGGATAAC TTAGTGTGTA GGGGGCCAGT GAGTTGGAGA TGAAGCGTA GGGAGTCGAA
5041 GGTGTCTTTT TGCGCCGAGT CAGTTCCTGG GTGGGGGCCA CAAGATCGGA TGAGCCAGTT TATCAATCCG
5111 GGGGTGCCAG CTGATCCATG GAGTGCAGGG TCTGCAAAAT ATCTCAAGCA CTGATTGATC TTAGGTTTTA
5181 CAATAGTGAT GTTACCCAG GAACAATTG GGAAGGTCA GAATCTTGTA GCCTGTAGCT GCATGACTCC
5251 TAAACATAA TTTCTTTTTT GTTTTTTTTT TTTATTTTTT GAGACAGGGT CTCACCTCTG CACCTAGGCT
5321 GAGTGCAGT GGTGCAATCA CAGCTCACTG CAGCCCTAG AGCGCCGCCC ACCGCGGTGG AGCTCCAATT
5391 CGCCCTATAG TGAGTCGTAT TACAATTAC TGCCCGTCTG TTTACAACGT CGTGACTGGG AAAACCCCTG
5461 CGTTACCCAA CTTAATCGCC TTGCAGCACA TCCCCCTTTC GCCAGCTGGC GTAATAGCGA AGAGGCCCGC
5531 ACCGATCGCC CTTCCCAACA GTTGCAGCAG CTGAATGGCG AATGGCGCGA AATTGTAAAC GTTAATATTT
5601 TGTTAAATTT CGCGTTAAAT TTTTGTAAA TCAGCTCATT TTTTAACCAA TAGGCCGAAA TCGGCAAAAT
5671 CCCTTATAA TCAAAAGAAT AGACCGAGAT AGGCTTGAGT GTTGTTCAG TTTGGAACAA GAGTCCACTA
5741 TTAAGAAACG TGGACTCCAA CGTCAAAGGG CGAAAAACCG TCTATCAGGG CGATGGCCCA CTACGTGAAC
5811 CATCACCTTA ATCAAGTTTT TTGGGGTCGA GGTGCCGTAA AGCACTAAAT CGGAACCCTA AAGGGAGCCC
5881 CCGATTTAGA GCTTGACGGG GAAAGCCGGC GAACTGTGGC AGAAAGGAAG GGAAGAAAGC GAAAGGAGCG
5951 GGCGCTAGGG CGCTGGCAAG TGTAGCGGTC ACGCTGCGCG TAACCACCCAC ACCGCGCGCG CTTAATGCCG
6021 CGCTACAGGG CGCGTCCCAG GTGGCACTTT TCGGGGAAAT GTGCGCGGAA CCCCTATTTG TTTATTTTTT
6091 TAAATACATT CAAATATGTA TCCGCTCATG AGACAATAAC CCTGATAAAT GCTTCAATAA TATTGAAAAA
6161 GGAAGAGTAT GAGTATTCAG CATTTCCGTG TCGCCCTTAT TCCCTTTTTT GCGGCATTTT GCCTTCTCTG
6231 TTTTGTCTAC CCAGAAACGC TGGTGAAGT AAAAGATGCT GAAGATCAGT TGGGTGCACG AGTGGGTTAC
6301 ATCGAAGTGG ATCTCAACAG CGGTAAAGAT CTTGAGAGTT TTGCCCCGGA AGAACGTTTT CCAATGATGA
6371 GCACTTTTAA AGTTCGTCTA TGTGGCGCGG TATTATCCCG TATTGACGCC GGGCAAGAGC AACTCGGTGC
6441 CGGCATACAC TATTCTCAGA ATGACTTGGT TGAGTACTCA CCAGTCACAG AAAGCATCT TACGGATGGC
6511 ATGACAGTAA GAGAATTATG CAGTGCTGCC ATAACCATGA GTGATAACAC TCGGCGCAAC TTACTTCTGA
6581 CAACGATCGG AGGACCGAAG GAGCTAACCG CTTTTTTGCA CAACATGGGG GATCATGTAA CTGCGCTTGA
6651 TCGTGGGAA CCGGAGCTGA ATGAAGCCAT ACCAAACGAC GAGCGTGACA CCAGATGCCC TGTAGCAATG

```

6721 GCAACAACGT TCGGCAAACT ATTAACCTGGC GAACTACTTA CTCTAGCTTC CCGGCAACAA TTAATAGACT  
6791 GGATGGAGGC GGATAAAGTT GCAGGACCAC TTCTGGGCTC GGCCTTCCG GCTGGCTGGT TTATTGCTGA  
6861 TAAATCTGGA GCCCGTGAGC GTGGGTCTCG CGGTATCATT GCAGCACTGG GGCAGATGG TAAGCCCTCC  
6931 CGTATCGTAG TTATCTACAC GACGGGGAGT CAGGCAACTA TGGATGAACG AAATAGACAG ATCCCTGAGA  
7001 TAGGTGCCCTC ACTGATTAAAG CATTTGGTAAC TGTACAGCCA AGTTTACTCA TATATACTTT AGATTGATTT  
7071 AAAACTTCAT TTTTAATTIA AAAGGATCTA GGTGAAGATC CTTTTIGATA ATCTCATGAC CAAAATCCCT  
7141 TAACGTGAGT TTTCTGTTCA CTGAGCGTCA GACCCCGTAG AAAAGATCAA AGGATCTTCT TGAGATCCCT  
7211 TTTTCTGCG CGTAATCTGC TGCTTGCAAA CAAAAAACC ACCGCTACCA GCGGTGGTTT GTTTGCCGGA  
7281 TCAAGAGCTA CCAACTCTTT TTCCGAAGGT AACTGGCTTC AGCAGAGCGC AGATACCAAA TACTGTCTTT  
7351 CTAGTGTAGC CGTAGTTAGG CCACCACTTC AAGAATCTG TAGCACCGCC TACATACCTC GCTCTGCTAA  
7421 TCCTGTTACC AGTGGCTGCT GCCAGTGGCG ATAAGTCGTG TCTTACCGGG TTGGACTCAA GACGATAGTT  
7491 ACCGGATAAG GCGCAGCGGT CGGGCTGAAC GGGGGGTTCC TGACACAGC CCAGCTTGA CCGAACGACC  
7561 TACACCGAAC TGAGATACCT ACAGCGTGAG CTATGAGAAA GCGCCACGCT TCCCGAAGGG AGAAAGGCGG  
7631 ACAGGTATCC GGTAAAGCGC AGGGTCGGAA CAGGAGAGCG CACGAGGGAG CTTCCAGGGG GAAACGCTG  
7701 GTATCTTTAT AGTCTCTGTC GGTTCGCCA CCTCTGACTT GAGCGTCGAT TTTTGTGATG CTCGTCAGGG  
7771 GGGCGGAGCC TATGAAAAA CGCCAGCAAC GCGGCTTTT TACGGTTCTT GGCCTTTTTC TGGCTTTTTC  
7841 CTCACATGTT CTTTCCCTGG TTATCCCTCG ATTCTGTGGA TAACCGTATT ACCGCTTTTG AGTGAGCTGA  
7911 TACCGCTGCG CGCAGCCGAA CGACCGAGCG CAGCGAGTCA GTGAGCGAGG AAGCGGAAGA GCGCCCAATA  
7981 CGCAAACCGC CTCCTCCCGC GCGTTGGCCG ATTCAATTAAT GCAGCTGGCA CGACAGGTTT CCGGACTGGA  
8051 AAGCGGGCAG TGAGCGCAAC GCAATTAATG TGAGTTAGCT CACTCATTAG GCACCCAGG CTTTACACTT  
8121 TATGCTTCCG GCTCGTATGT TGTGTGGAAT TGTAGCGGA TAACAATTTC ACAACGGAAA CAGTATGAC  
8191 CATGATTACG CCAAGCTCGG AATTAACCTT CACTAAAGG AAAAAAGCT GCTGCAGGCT CCCTAACTGC  
8261 CAAGCCCCAC AGTGTGCCCT GAGGCTGCCC CTTCCTTCTA GCGGCTGCCC CCACTCGGCT TTGCTTTCCC  
8331 TAGTTTCAGT TACTTGGCTT CAGCCAAGGT CTGAAACTAG GTGCGCACAG AGCGGTAAGA CTGCGAGAGA  
8401 AAGAGACCAG CTTTACAGGG GGTTTATCAC AGTGCACTCT GACAGTCTGC AGCCTCACAG GGGGTTTATC  
8471 ACATTGCAAC CTGACAGTCC TCAGCCTCAC AGGGGGTTTA TCACAGTGCA CCCTTACAAT CATTCATT  
8541 GATTACAAT TTTTCTAGTC TCTACTGTGC CTAACCTGTA AGTTAAATTT GATCAGAGGT GTGTCCAG  
8611 AGGGGAAAAC AGTATATACA GGGTTCACTA CTATCGCATT TCAGGCTTCC ACCTGGGTCT TGGAATGTGT  
8681 CCCCCGAGGG GTGATGACTA CCTCAGTTGG ATCTCCACAG GTCACAGTGA CACAAGATAA CCAAGACACC  
8751 TCCCAAGGCT ACCACAATGG GCGGCTTCC ACGTGCAAT GCGCGGAGGA ACTGCCATGT CGGAGGTGCA  
8821 AGCACACCTG CGCATCAGAG TCCTTGCTGT GAGGGGAGGG ACCAGCGCAG CTTCCAGCCA TCCACCTGAT  
8891 GAACAGAAC TAGGGAAAGC CCCAGTTCTA CTTACACCAG GAAGGC

32/43

FIGURE 12

```

mBCwCN frag  -----C- --AC--G--
m2BCwCN frag  -----C- --G--G--
BC/HXB2       -----
BC/NL43       -----
#1            .....
              CGCGCACGGC AAGAGGCGAG GGGCGGCGAC TGGTGAGTAC GCCAAAAATT

```

```

mBCwCN frag  -----C- C-----
m2BCwCN frag  -----
BC/HXB2       -----T-----
BC/NL43       -----G-----
#51           .....
              TTGACTAGCG GAGGCTAGAA GGAGAGAGAT GGGTGCGAGA GCGTCAGTAT

```


```

mBCwCN frag  -----
m2BCwCN frag  -----
BC/HXB2       -----
BC/NL43       -----AA-----
#101         .....
              TAAGCGGGGG AGAATTAGAT CG

```


33/43

FIGURE 13




34/43

FIGURE 14




35/43

FIGURE 15A



36/43


FIGURE 15B






37/43

FIGURE 15C



38/43

FIGURE 15D



39/43

FIGURE 16




FIGURE 17

1 CCTGCCCAAT GCATAOFTTG TATCCATATC ATAATATGTA CATTATATT GOCTCATGTC CAACATPACC  
 71 GCCATGTTGA CATTGATTAT TGA CTAGTTA TTAATAGTAA TCAATTACGG GTTCATPAGT TCATAGCOCA  
 141 TATATGGAGT TCCGCTTAC ATAACTTACG GTAATGCGC CCGCTGCTG ACGCCCAAC GACCCCGCC  
 211 CATTGACGTC AATAATGACG TATGTTCCCA TAGTAAGCC AATAGGCACT TTCATTTGAC GTCAATGGGT  
 281 GGATATTTTA CGGTAAACTG CCCACTTGGC AOTACATCAA GNTATCATA TOCCAAGTAC GCCCCCTATT  
 351 GACOTCAATG ACGGTAAATG GCGCGCTGG CATTATGCC AGTACATGAC CTTATGGUAC TTTCCTACTT  
 421 GGCAGTACAT CTAGCTATTA GTATCGCTA TTACCATGCT GATGCGGTTT TGGCAGTACA TCAATGGGGG  
 491 TGGATAGCGG TTGACTCAC GGGGATTTCC AAGTCTCCAC CCCATGACG TCAATGGGAG TTCTGTTTGG  
 561 CACCAAAATC AACGGGACTT TCCAAATGT GGTAACTACT CGGCTCAAT GACGCAATG GCGGTTAGGC  
 631 GTGTACGGTG GGAGGTCTAT ATAAGCAGAG CTGCTTACTT GAACCTCAG ATCGCTGGA GACGCTATCC

BssHII (752)

SacII (746)

701 ACGCTGTTTT GACCTCCATA GAAGACACCG GGACCGATCC AGCCTCCGCG GGCGCGCTAA GTATGGGATG  
 771 TCTTGGGAAT CAGCTGCTTA TCGCCATCTT GCTTTTAAAT GTCTATGGGA TCATATGTAC TCTATATGTC  
 3> L G N Q L L I A I L L L S V Y G I Y C T L Y V  
 841 ACAOTCTTTT ATGGTGTACC AGCTTGCAGG AATGGACAA TTCCCTCTT TTGTGCAACC AAGATAGGG  
 27> T V F Y G V P A W R N A T I P L F C A T K N R  
 911 ATACTGGGG AACAACTCAG TGCTACCGAG ATAATGGTGA TTAATCAGAA GTGGCCCTTA ATATACAGA  
 56> D T W G T T Q C L P D N G D Y S E V A L N V T E  
 981 AAGCTTTGAT CCGTGAATA ATACACTCAC AGAACACGCA ATAGAGGATG TATGGCAACT CTTTGAGACC  
 73> S F D A W N N T V T E Q A I E D V W Q L F E T  
 1051 TCATTAAGC CTGTGTAAA ATTATCCCA TTATGCCATTA CTATGAGATG CAATAAAGT GACACAGATA  
 97> S I K P C V K L S P L C I I M R C N K S E T D  
 1121 GATGGGGATT GACAAATCA ATACAAACA CAGCATCAAC AACATCAACG ACAGCATCAG CAAAGTAGA  
 120> R W G L T K S I T T A S T T S T T A S A K V D  
 1191 CATGGTCAAT GAGACTAGTT CTATATAGC CCGAGTAAAT TGCAACAGCT TGAACAAGA GCAATGATA  
 143> M V N E T S S C I A Q D N C T G L E Q E Q M I  
 1261 AGCTGTAAAT TCAACATGAC AGGTTTAAA AGAGCAAGA AAAAAGATA CAATGAACT TGTACACTG  
 167> S C K F N M T G L K R D K K K E Y N E T W Y S  
 1331 CAGATTTCCT ATGTGAACAA GCGAATAACA CTGTAAATGA AAGTAGATGT TACATGAACC ACTGTAAACAC  
 190> A D L V C E Q G N N T G N E S R C Y M N H C N T  
 1401 TTCTGTATC CAAGAGTCTT GTGACAAACA TTATTTGGAT GCTATTAGAT TTAGGTATG TGCACCTCCA  
 213> S V I Q E S C D K H Y W D A I R F R Y C A P P  
 1471 GGTATGCTT TCTTAGATG TATGACACA AATTATTGAG GCTTTATGCC TAAATGTCT AAGGTGGTGG  
 237> G Y A L L R C N D T N Y S G F M P K C S K V V  
 1541 TCTCTTCATG CACAGGATG ATGAGACAC AGACTTCTAC TTGGTTTGGC TTAAATGGAA CTAGACGAGA  
 260> V S S C T R M M E T Q T S T W F G F N G T R A E  
 1611 AAATAGAACT TATATTTACT GGCATGGTAG GGATAATAGG ACTATAATTA GTTAAATAA GTATTATAAT  
 283> N R T Y I Y W H G R D N R T I I S L N K Y Y N  
 1681 CTACAATGA AATGTAGAAG ACCAGGAAT AAGACAGTTT TACCAGTAC CATATGTCT GCATTTGTTT  
 307> L T M K C R R P G N K T V L P V T I M S G L V  
 1751 TCCATTCACA ACCAATCAAT GATAGCCAA AGCAGGCATG GTGTTGGTTT GCGGAAAT GCGAGGATC  
 330> F H S Q P I N D R P K Q A W C W F G G K W K D A  
 1821 AATAAAGAG GTGAAGCAGA CCATTGTCAA ACATCCAGG TATACTGGA CTACCAATAC TGATAAATC  
 353> I K E V K Q T I V K H P R Y T G T N N T D K I  
 1891 AATTTGACCG CTCCTGAGG AGGAGATCCG GAAGTTACCT TCATGTGAC AAAATTGAGA GACAGTATC  
 377> N L T A P G G G D P E V T F M W T N C R G E F

1961 TCTACTGTAA AATGAATTGG TTCTTAAATT GGGTAGAAGA TAGGAATACA GCTAACCAGA AOCCTAAAGGA  
 400 L Y C K M N W F L N W V E D R N T A N Q K P K E  
 2031 ACAGCATAAA AGGAATTACG TGOCATGTCA TATTAGACAA ATAATCAACA CTTGGCATAA AGTAGGCAAA  
 423 Q H K R N Y V P C H I R Q I I N T W H K V G K  
 2101 AATGTTTATT TGCTTCCAAG AGAGGGAGAC CTCACGTGTA ACTTCACAGT GACCAGTCTC ATAGCAAAACA  
 447 N V Y L P P R E G D L T C N S T V T S L I A N  
 2171 TAGATTGGAT TGATGGAAAC CAACTAATA TCACCATGAG TGCAGAGGTG GCAGAACTGT ATCGATTGGA  
 470 I D W I D G N Q T N I T M S A E V A E L Y R L E  
 2241 APTGGGAGAT TATAAATTAG TAGAGATCAC TCCANTTGGC TTGGGXXCCA CAGATGTGAA GAGGTACACT  
 493 L G D Y K L V E I T P I G L A P T D V K R Y T  
 2311 ACTGGTGGCA CCTCAAGAAA TAAAGAGGGG GTCTTTTGTG TAGGGTTCTT GGGTTTTCTC GCAACGGCAG  
 517 T G G T S R N K R G V F V L G F L G F L A T A  
 2381 GTTCTGCAAT GGGAGCCGCC AGCCTGACCC TCAGGGCACA GTCCCGAACT TTATTGGCTG GGAATAGTCA  
 540 G S A M G A A S L T L T A Q S R T L L A G I V Q  
 2451 ACAGCAGCAA CAGCTGTGCG AGCTGTGCAA GAGACAACA GAATTTTGAU GACTGACCGT CTCGGGACAA  
 563 Q Q Q Q L L D V V K R Q Q E L L R L T V W G T  
 2521 AAGAACCTCC AGACTAGGGT CACTGCCATC GAGAGTACT TAAAGGACCA GCGCGAGCTG AATGCTTGGG  
 587 K N L Q T R V T A I E K Y L K D Q A Q L N A W  
 2591 GATGTGCGTT TAGCAAGTC TGCCACACTA CTGTACCATG GCCAATCCA AGCTAACAC CAAAGTGGAA  
 610 G C A F R Q V C H T T V P W P N A S L T P K W N  
 2661 CAATGAGACT TGGCAAGAGT GGGAGCGAAA GGTGACITC TTGGAGAGAA ATATAACAGC CCTCTACAG  
 633 N E T W Q E W E R K V D F L E E N I T A L L E  
 2731 GAGGCACAAA TTCAACAAGA GAAGACATG TATGAATTAC AAAAGTTGAA TAGCTGGGAT GTGTTTGGCA  
 657 E A Q I Q Q E K N M Y E L Q K L N S W D V F G  
 2801 APTGGTTTGA CCTGCTTCTT TGATATAAGT ATATACAATA TGGAGTTTAT ATAGTTGTAG GAGTAATACT  
 680 N W F D L A S W I K Y I Q Y G V Y I V V G V I L  
 2871 GTTAAAGATA GTGATCTATA TAGTACAAAT GCTAGCTAAG TTAAGGCAGG GGTATAGGCC AGTGTCTCT  
 703 L R I V I Y I V Q M L A K L R Q G Y R P V F S  
 2941 TCCCCACCTT CTTATTTCOA GCAGACCAT ATCCAACAGG ACCCGGCACT GCCAACCAGA GAAGGCAGAG  
 727 S P P S Y F Q Q T H I Q Q D P A L P T R E G K  
 3011 AAAGAGACCG TGAGAGAGGC GGTGGCAACA GCTTCTGGCC TTGGCAGATA GAATATATCC ACTTCTTTAT  
 750 E R D G G E G G G N S S W P W Q I E Y I H F L I  
 3081 TGTGAGCTT APTGAGCTT TGACTTGGCT ATTCACTAAC TGTAGGACTT TGTATATAG ACTATACAG  
 773 R Q L I R L L T W L F S N C R T L L S R V Y Q  
 3151 ATCCTCCAAC CAATACTCCA GAGGCTCTCT GCGACCTTAC AGAGGATTCT AGAGTCTCT AGGACTGAC  
 797 I L Q P I L Q R L S A T L Q R I R E V L R T E  
 3221 TGACCTACTT ACAATATGGG TGGAGCTATT TCCATGAGGC GGTCCAGGCC GTCTGGAGAT CTGCGGACCG  
 820 L T Y L Q Y G W S Y F H E A V Q A V W R S G D R  
 3291 ACAGAGACTC TTGGGGGCCG GTGGGGGAGC TTATGGGAGA CTCTTAGAGA CTCTTGGGGG CGCCTTGGGA  
 843 Q R L L R A R G E T Y G R L L E T L A G A W G  
 3361 GACTTATGGG AGACTCTTAG GAGAGGAGAG GTGGAGAGAT GATACTCGAG GTGGAAGATG GATACTCGCC  
 867 D L W E T L R R G E V E D G Y S R W K M D T R  
 3431 AAATOCATCC CAGAGGAGT TAGACAAGGG CTTGAGCTCA CTCCTTGTG ACCAGGAGGA TTAGACAAGG  
 890 Q I H P Q E D

**BamHI (3539)**

3501 GCTTGAAGCTC ACTCTCTTGT GAGGACAGCA GGGACAGAGg atccactagt tctagaCTCG AGGGGGGGGCC  
 3571 GGTACGAGC GTTAGCTAG CTAGAGACCA CCTCCCTTGC GAGCTAAGCT GGACAGCCAA TGACGGGTAA  
 3641 GAGAGTGACA TTTTTCATTA ACCTAAGACA GGAGGGCCGT CAGAGCTACT GCTAATCCA AAGACGGGTA  
 3711 AAGTGATAA AAATGTATCA CTCCAACCTA AGACAGGGCG AGCTTCCGAG GGATTTGTG TCTGTTTTAT  
 3781 ATATATTTAA AAGGGTGACC TGTCCGGAGC CGTGTGCCC GGATGATGTC TTGGTCTAGA CTCGAGGGGG  
 3851 GGCCCCGTAC GATCCAGATC TGTGTGCGCT TCTAGTGGCC AGCCATCTCT TGTPTGCTCC TCCCGGCTC  
 3921 CTTCTTGGAC CTTGGAGGT GCCACTCCCA CTGCTCTTTC CTAAATTAAT GAGGAATTG CATCGCATG

3991 TCTTAGTAGG TGTATTTCTA TTCTAGGGG TGGTATGAG CAGCAGACCA AGGAGAGGA TTGAGAGAC

### KpnI (4097)

4061 AATAGCAGGC ATGCTGGGGA TGGGTGGGC TGTATAGGTA CCCAGGTGCT GAAGAATTGA CCGGTTCCT

4131 CTTAGGCCAG AAAGAGCCAG GCAATGCCG TTCTATGTA CACACCTGCT CCAAGCCCTT GTTCTTAGT

4201 TCCAGCCCCA CTCATAGGAG ACTATAGCT CAGGAGGCTT CCGCTTCAA TCCAGCCG TAAAGTACTT

4271 GAGCGGTCT CTCCCTCCTT CATCAACCA CCAAGCCAA CTTAGCTTCC AAGAGTGGGA AGAATTAA

4341 CCAAGATGAG CTATTAGTG CAGAGGAGA GAAATGCC TCAACATGTC AGGAGTAAT GAGAGAAAT

4411 ATAGAATTTC TTCCGCTTCC TCGCTCACTG ACTCGCTGCG CTGGGTGCTT CCGCTGGCGC GAGCGGTATC

4481 AGCTCACTCA AAGGCGGTAA TACGGTTATC CACAGAATCA GGGGATAACG CAGGAAAGAA CATGTGACCA

4551 AAAGGCCAGC AAAAGGCCAG GAACCGTAAA AAGGCGCGCT TGCTGGCGTT TTTCATAGG CTCGCCCCC

4621 CTGACGAGCA TCACAAAAT CGACGCTCAA GTCAGAGGTG GCGAAACCG ACAGGACTAT AAAGATAOCA

4691 GCGTTTCCC OCTGGAAGCT CCTCGTGGC CTCTCTGTT CCGACCTGC CGCTTACGG ATACCTGTCC

4761 GCCTTTCTCC CTTCGGGAAG CGTGGCGCTT TCTCAATGCT CACGCTGTAG GTATCTCAGT TCGGTGTAGG

4831 TCGTTCGCTC CAAGCTGGGC TGTGTGCAG AACCCCCGT TCAGCCGAC CGCTGGCGCT TATCCGGTAA

4901 CTATCGTCTT GAGTCCAACC CGGTAAGACA CGACTTATCG CCACCTGGCAG CAGCCACTGG TAACAGGATT

4971 AGCAGAGCGA GGTATGTAGG CGGTGCTACA GAGTCTTGA AGTGGTGGCC TAACTACGC TACACTAGAA

5041 GGACAGTATT TGGTATCTGC GCTCTGCTGA AGCCAGTTAC CTTCGGAAAA AGAGTTGGTA GCTCTGTATC

5111 CCGCAAACAA ACCACGCTG GTAGCGGTGG TTTTGTGTT TGCAAGCAGC AGATTACGG CAGAAAAAA

5181 GGATCTCAAG AAGATCCTTT GATCTTTTCT ACGGGCTCG ACCTCAGTG GAACGAAAC TCACGTTAAG

5251 GGATTTTGGT CATGAGATTA TCAAAAAGGA TCTTCACCTA GATCCTTTTA AATTAAAAAT GAAGTTTAA

5321 ATCAATCTAA AGTATATATG AGTAAACTTG GTCTGACAGT TACCAATGCT TAATCAGTGA GGCACCTATC

5391 TCAGCGATCT GTCTATTTCC TTTATCCATA GTTGCTGAC TCCGGGGGGG GGGGGCGCT AGGTCTGCT

5461 CGTGAAGAAG GTGTGCTGA CTCATACCAG GCCTGAATCG CCCCATCATC CAGCCAGAAA GTGAGGGAGC

5531 CACGGTTGAT GAGAGCTTTG TTGTAGGTGG ACCAGTTGGT GATTTTGAAC TTTTGTCTTG CCACGGAACG

5601 GTCTGCGTGG TCGGAAGAAT GCGTGAATCT ATCCTTCAAC TCAGCAAAAG TTCGATTAT TCAACAAAGC

5671 CGCGCTCCCG TCAAGTCAGC GTAATGCTCT GCCAGTGTTA CAACCAATTA ACCAATCTCT ATTAGAAAA

2714 F F

5741 CTCATCGAGC ATCAAAATGAA ACTGCAATTT ATTATATCA GGATTATCAA TACCATATTT TTGAAAAAGC

2694 E D L M L H F Q L K N M D P N D I G Y K Q F L

5811 CGTTTCTGTA ATGAAGGAGA AAATCACCAG AGCGAGTTCC ATAGGATGGC AAGATCCTGG TATCGGTCTG

2454 R K Q L S P S F E G L C N W L I A L D Q Y R D A

5881 CGATTCCGAC TGTCCAACA TCAATACAAC CTATTAATTT CCCCCTGTCA AAAATAAGGT TATCAAGTGA

2224 I G V R G V D I C G I L K G E D F I L N D L S

5951 GAAATCACA TGAATGACGA CTGAATCCGG TGAGAATGGC AAAAGCTTAT GCATTTCTTT CCAGACTTGT

1994 F D G H T V V S D P S F P L L K H M E K W V Q

6021 TCAACAGGCC AGCCATTACG CTCGTATCA AAATCACTCG CATCAACCAA ACGTTATTC ATTCTGATT

1754 E V P W G N R E D D F D S A D V L G N N M R S Q

6091 GCGCTGAGC GAGACGAAAT ACGCGATCG TGTAAAAGG ACAATTACAA ACAGGAATCG AATGCAACCG

1524 A Q A L R F V R D S N F P C N C V P I S H L R

6161 GCGCAGGAA ACTGCCAGCG CATCAACAAT ATTTTACCTT GAATCAGGAT ATTCTTCTAA TACCTGGAAT

1294 R L F V A L A D V I N E G S D P Y E E L V Q F

6231 GCTGTTTTC CCGGGATCGC AGTGGTGAAT AACCATGCAT CATCAGGAGT ACGGATAAAA TGCTTGATGG

1054 A T K G P I A T T L L W A D D P T R I F H K I T

6301 TCGGAAGAGG CATAAATTC GTCAGCCAGT TTAGTCTGAC CATCTCATCT GTAACATCAT TGGCAACGCT

824 P L P M F E T L W N L R V M E D T V D N A V S

6371 ACCTTTCCA TGTTTCAGAA ACAACTCTGG CGCATCGGC TTCCCATACA ATCGATAGAT TGTGCACT

594 G K G H K L F L E P A D P K G Y L R Y I T A G

6441 GATTACCGCA CATATCGCG AGCCCATTTA TACCATATA AATCAGCATC CATGTTGGAA TTATATCGC

354 S Q G V N D R A W K Y G Y L D A D M N S N L R P

6511 GCCTCAGCA AGACGTTTCC CGTTGAATAT GGCTCATAAC ACCCCTTGTA TTAATGTTA TGTAAAGCAG

124 R S C S T E R Q I H S M

6581 CAGTTTATT GTTCATGATG ATATATTTT ATCTGTGCA ATGTAACATC ACAGATTTTG AGACACAACG

43/43

6651 TGGCTTTCCC CCCCCCCCCA TTATTGAAGC ATTTATCAGG GTTATTGTCT CATGAGCGGA TACATATTTG  
6721 AATGTATTTA GAAAAATAAA CAAATAGGGG TTCCGCGCAC ATTTCCCCGA AAAGTGCCAC CTGACGTCTA  
6791 AGAAACCATT ATTATCATGA CATTAACTTA TAAAAATAGG CGTATCACGA GGCCCTTTTG TCTCGCGCGT  
6861 TTCGGTGATG ACGGTGAAAA CCTCTGACAC ATGCAGCTCC CGGAGACGGT CACAGCTTGT CTGTAAAGCG  
6931 ATGCCGGGAG CAGACAAGCC CGTCAGGGCG CGTCAGCGGG TGTTCGGGGG TGTTCGGGGCT GGCTTAACTA  
7001 TGCGGCATCA GAGCAGATTG TACTGAGAGT GCACCATATG CGGTGTGAAA TACCGCACAG ATGCGTAAGG  
7071 AGAAAATACC GCATCAGATT GGCTATTGG